

Figure 1-3f. Sensitive Biological Resources Between Shasta Dam and Red Bluff Diversion Dam

This page left blank intentionally.




Figure 1-3g. Sensitive Biological Resources Between Shasta Dam and Red Bluff Diversion Dam

This page left blank intentionally.




Figure 1-3h. Sensitive Biological Resources Between Shasta Dam and Red Bluff Diversion Dam

This page left blank intentionally.




Figure 1-3i. Sensitive Biological Resources Between Shasta Dam and Red Bluff Diversion Dam

This page left blank intentionally.




Figure 1-3j. Sensitive Biological Resources Between Shasta Dam and Red Bluff Diversion Dam

This page left blank intentionally.

Valley Oak Woodland This habitat type consists of an open savanna of valley oak (*Quercus lobata*) trees and an annual grassland understory. Valley oak is typically the only tree species present and shrubs are generally absent except for occasional poison oak. Canopy cover rarely exceeds 30–40 percent in valley oak woodland. This community occupies the highest portions of the floodplain terrace where flooding is infrequent and shallow. Valley oak woodland is a community in the MSCS habitat type “valley/foothill woodland and forest.”

The oaks that dominate the tree layer of oak savannas and woodlands are long-lived trees that are resilient to damage; their stems often survive fire, and when their stems are killed by fire or are cut down, basal sprouts often grow into new stems. (Valley oak also tolerates inundation during winter before it has leafed out.)

Nonetheless, there are concerns regarding the status and ongoing trends of tree canopies of blue oak- and valley oak-dominated savannas and woodlands (Tyler et al. 2006). When canopy oaks die, their replacement through the growth of saplings or the remaining canopy trees is necessary to maintain the tree canopy. Currently, saplings are absent or at low densities on many sites, and thus, their recruitment into the canopy is very infrequent. The infrequent recruitment of saplings appears to be a consequence of climate, interactions with herbivores, competition for resources with herbaceous plants, and the reproductive biology of the oaks themselves.

Riparian Communities Riparian communities in the primary study area are subject to regulation by DFG under Section 1602 of the California Fish and Game Code because they are associated with streambanks. They are identified as sensitive natural communities by DFG because of their declining status statewide and because of the important habitat values they provide to both common and special-status plant and animal species. These habitat types are tracked in the CNDDB. In addition, areas containing riparian habitat may be subject to USACE jurisdiction under Section 404 of the CWA, if they meet the three wetland criteria described below under “Regulatory Setting.”

California’s riparian and wetland communities have experienced the most extensive reductions in their acreage, and in the Sacramento Valley approximately 90 percent of riparian vegetation has been converted to agriculture or development, and the remainder substantially altered by dams, diversions, gravel mining, grazing practices, and invasive species (Hunter et al. 1999; CALFED 2000b). Consequently, riparian and wetland communities are considered sensitive.

Much of the vast riparian habitat that once existed along the Sacramento River has been eliminated by agricultural clearing, flood control projects, and urbanization. Historically, belts of riparian forest more than 5 miles wide occurred along the Sacramento River (Jepson 1893; Thompson 1961, cited in

Hunter et al. 1999). Only narrow remnants of these riparian forests remain in the Sacramento River Valley. In the primary study area, much of the

Sacramento River from Shasta Dam to Redding is deeply entrenched in bedrock, which precludes development of extensive areas of riparian vegetation. The river corridor between Redding and Red Bluff, however, still maintains extensive areas of riparian vegetation.

Riparian communities present within the floodplain of the Sacramento River, within the study area, include blackberry scrub, Great Valley willow scrub, Great Valley cottonwood riparian forest, Great Valley mixed riparian forest, and Great Valley valley oak riparian forest. Willow and blackberry scrub and cottonwood- and willow-dominated riparian communities are present along active channels and on the lower flood terraces whereas valley oak-dominated communities occur on higher flood terraces. These communities are all in the MSCS habitat type “valley/foothill riparian,” and are described below.

Blackberry Scrub Blackberry scrub is dominated by Himalayan blackberry (*Rubus discolor*), a species that is listed as invasive by the California Invasive Plant Council (Cal IPC). Cover of Himalayan blackberry is extremely dense in this community, leaving little opportunity for the establishment of native tree seedlings or shrubs beneath its canopy. Herbaceous cover is also very sparse. Scattered individual trees and shrubs may be interspersed through the blackberry scrub community. Himalayan blackberry generally establishes in gaps created by natural or human disturbances. Although Himalayan blackberry is an invasive species, this community does provide nesting habitat for some bird species and may be regulated under Section 1602 of the California Fish and Game Code when located within the bed, channel, or bank of a stream and may be subject to USACE jurisdiction under Section 404 of the CWA.

Great Valley Willow Scrub Great Valley willow scrub is a deciduous broadleaved community with open to dense cover of shrubby willows. This community type may be dominated by a single species of willow or by a mixture of willow species. Dense stands have very little understory while more open stands have herbaceous understories, usually dominated by grasses characteristic of the annual grassland community. Characteristic plant species include sandbar willow (*Salix exigua*), arroyo willow (*S. lasiolepis*), shining willow (*S. lucida*), and California wild rose (*Rosa californica*). This community occupies point bars and narrow corridors along the active river channel that are repeatedly disturbed by high flows.

Great Valley Cottonwood Riparian Forest Great Valley cottonwood riparian forest is a deciduous broadleaved forest community with a dense tree canopy dominated by Fremont cottonwood (*Populus fremontii*) and often including a high abundance of black willow (*Salix gooddingii*). This community also has a dense understory of seedlings, saplings, and sprouts of the canopy dominants and shade-tolerant species including California box elder

(*Acer negundo*) and Oregon ash (*Fraxinus latifolia*). Characteristic shrub species include California buttonbush (*Cephalanthus occidentalis* var. *californicus*) and willows (*Salix* spp.). Lianas such as California grape (*Vitis californica*) are typically present as well. This community occupies lower floodplain terraces that are flooded annually.

Great Valley Mixed Riparian Forest Great Valley mixed riparian forest is a deciduous broadleaved forest community with a moderately dense to dense tree canopy that typically includes several species as codominates. Shrubs and lianas are also typically present. Mixed riparian forest is typically dominated by sycamore (*Platanus racemosa*) and valley oak (*Quercus lobata*) with Fremont cottonwood, white alder (*Alnus rhombifolia*), willow, and Oregon ash also occurring frequently. Common shrub species in this community type include blue elderberry (*Sambucus mexicana*), California buttonbush, spicebush (*Calycanthus occidentalis*), and Himalayan blackberry. The herbaceous understory consists primarily of annual grasses and forbs similar to those found in the annual grassland communities but with a higher proportion of shade-tolerant species such as miner's lettuce (*Claytonia parviflora*), common bedstraw (*Galium aparine*), bur-chervil (*Anthriscus caucalis*), and meadow nemophila (*Nemophila pedunculata*). At stream edges, the herbaceous understory of this community is characterized by species that grow entirely or partially in water, such as tall flatsedge (*Cyperus eragrostis*), common tule (*Scirpus acutus*), cattail (*Typha* spp.), sedges (*Carex* spp.), deergrass (*Muhlenbergia rigens*), and common monkeyflower (*Mimulus guttatus*). This community occupies intermediate flood terraces that are subject to occasional high-flow disturbance.

Great Valley Valley Oak Riparian Forest Great Valley valley oak riparian forest is a deciduous broadleaved forest community with a closed canopy. This community type is similar to the Great Valley mixed riparian forest community described above but is clearly dominated by valley oak. Characteristic species include many of the same associates found in the Great Valley mixed riparian forest community type, but tree and shrub associates are more widely scattered. This community occupies upper floodplain terraces where flooding is infrequent but soil moisture is high.

More than 15 native species of deciduous trees and shrubs occur in the riparian forests, woodlands, and scrubs of the Central Valley and the Delta (Conard et al. 1977; Sawyer and Keeler-Wolf 1995; Vaghti and Greco 2007). Flow regime, disturbance, and species attributes determine the species composition and physical structure of this woody vegetation. Although flow regime influences the dispersal, establishment, growth, and survival of all the woody riparian species, Fremont's cottonwood (*Populus fremontii*) and the willow species (*Salix* sp.) particularly depend on specific hydrologic events for their recruitment. During seed release, flows must be high enough to disperse seed to surfaces where scouring by subsequent flows does not occur, yet not so high that seedlings desiccate after flows recede, and flows must recede gradually to

enable germination and seedling establishment while the substrate is still moist (Mahoney and Rood 1998).

Fremont's cottonwood and willow species are rapidly growing, shade intolerant and relatively short-lived (Burns and Honkala 1990, Vaghti and Greco 2007). Within 10–20 years, initially shrubby thickets have reached 10–40 feet in height. Other species, such as Oregon ash (*Fraxinus latifolia*) and valley oak (*Quercus lobata*), establish concurrently or subsequent to the willows and cottonwood, grow more slowly but are more tolerant of shade, and are longer-lived (Burns and Honkala 1990, Tu 2000). In the absence of frequent disturbance, these species enter the canopy, particularly after 50 years, as mortality of willows and cottonwood frees space. Conversely, frequent disturbance prevents the transition to mature mixed riparian or valley oak forests.

Riparian trees and shrubs are readily top-killed by fire (valley oak is somewhat more resistant to fire); however, most species will produce new shoots from the base of their previous trunks or from their roots. Important consequences of frequent fire can include shifts in species composition, more open vegetation, and an absence of larger trees (which are important habitat components for some species of wildlife).

None of the native woody species of the Central Valley's riparian areas germinate and establish seedlings underwater, and all tolerate inundation during the dormant season (i.e., late fall to early spring). Tolerance of inundation during the growing season varies among species. Most species can withstand a prolonged period of inundation if at least a portion of their foliage is above the water surface.

Wetland Communities Similar to riparian communities, much of the wetland habitat that once occurred in the Sacramento River Valley has been eliminated as a consequence of land use conversion to agriculture and urbanization. It is estimated that nearly 1.5 million acres of wetlands once occurred in the Central Valley. Today, approximately 123,000 acres remain. Wetland communities that are likely to occur in the primary study area between Shasta Dam and RBDD include freshwater marsh, freshwater seep, northern hardpan vernal pools, northern volcanic mudflow vernal pools, and other seasonal wetlands. Riparian communities in the primary study area are subject to regulation by DFG under Section 1602 of the California Fish and Game Code because they are associated with streambanks. They are identified as sensitive natural communities by DFG because of their declining status statewide and because of the important habitat values they provide to both common and special-status plant and animal species. These habitat types are tracked in the CNDDDB. In addition, areas containing riparian habitat may be subject to USACE jurisdiction under Section 404 of the CWA, if they meet the three wetland criteria described below under "Regulatory Setting."

Freshwater Marsh Freshwater marshes are herbaceous wetland plant communities that occur along rivers and lakes and are characterized by dense cover of perennial, emergent plant species. Marshes are typically perennial wetlands, but may dry out for short periods of time. Characteristic freshwater marsh species include common tule (*Scirpus acutus*), narrowleaf cattail (*Typha angustifolia*), broadleaf cattail (*Typha latifolia*), common reed (*Phragmites australis*), tall flatsedge, common spikerush (*Eleocharis macrostachya*), and sedges (*Carex* spp.). Freshwater marshes in the primary study area are communities in the MSCS habitat type “nontidal freshwater permanent emergent.”

In marsh vegetation, vegetation structure and species richness are strongly influenced by disturbance, changes in water levels, and the range of elevations present at a site (Keddy 2000). Disturbances, and water level drawdowns that expose previously submerged surfaces, provide opportunities for annuals, short-lived perennials, and other species to establish, which creates diversity in species composition and vegetation structure.

Although seedling establishment takes place on exposed surfaces, clonal growth allows many plants to subsequently occupy sites at lower elevations (i.e., in deeper water). The growth of emergent macrophytes is reduced, however, by submergence and by damage to their culms from wave action; thus vegetation dominated by emergent macrophytes is restricted to shallow water, typically less than 2 feet in depth (Coops et al. 1991; Coops et al. 1996).

At upper and lower elevations, wetlands frequently intergrade with adjacent vegetation. At upper elevations, emergent wetlands often intergrade with the woody vegetation of adjacent riparian areas; at lower elevations, wetlands often intergrade with aquatic vegetation, and a characteristic feature of this transitional zone is floating mats of plants such as water primrose (*Ludwigia* spp.) and knotgrass (*Paspalum distichum*).

Freshwater Seep Freshwater seep is a wetland plant community characterized by dense cover of perennial herb species usually dominated by rushes, sedges, and grasses. Freshwater seep communities occur on sites with permanently moist or wet soils resulting from daylighting groundwater. Species commonly observed in freshwater seeps in the area include rushes (*Juncus* spp.), sedges, flatsedges (*Cyperus* spp.), deergrass, cattail, bull thistle (*Cirsium vulgare*), blue-eyed grass (*Sisyrinchium bellum*), and willow (*Salix* spp.). The MSCS does not designate a habitat type that includes freshwater seeps.

Vernal Pools Vernal pools are seasonal wetlands that fill during winter rains and dry up in spring. They occur in undulating or mima mound (i.e., mound-intermound) topography where the soil or underlying rock has layers that are relatively impermeable to water. Vernal pools may be isolated from one another, but more often they are interconnected by swales or ephemeral

drainages in vernal pool complexes that may extend for hundreds of acres. Vernal pool complexes generally include water features.

Vernal pools are considered sensitive because they provide potential habitat for Federally listed species, including slender orcutt grass (*Orcuttia tenuis*) and vernal pool crustaceans; provide important ecological values and functions; and are likely considered waters of the State subject to jurisdiction of the Central Valley Regional Water Quality Control Board under the Porter-Cologne Act.

Vernal pool communities are included in the MSCS habitat type “natural seasonal wetland.” The two predominant types of vernal pool communities in the study area, northern hardpan vernal pools and northern volcanic mudflow vernal pools, are described below.

Northern Hardpan Vernal Pool This is an herbaceous plant community characterized by low-growing annual grasses and forbs adapted to live both on land and in water. In this type of vernal pool, the restrictive layer is composed of cemented hardpan soils that are very acidic. This type of vernal pool occurs on alluvial terraces with hummocky (i.e., mound and depression) topography. The pools form in depressions between mounds. Northern hardpan vernal pool communities are dominated by native plant species; characteristic species include Fremont’s goldfields (*Lasthenia fremonitii*), toothed downingia (*Downingia cuspidata*), Sacramento Valley pogogyne (*Pogogyne zizyphoroides*), stalked popcornflower (*Plagiobothrys stipitatus*), common blennosperma (*Blennosperma nanum*), pigmy-weed (*Crassula aquatica*), and water starwort (*Callitriche marginata*).

Northern Volcanic Mudflow Vernal Pool This is an herbaceous plant community characterized by an open mixture of very low-growing annual grasses and forbs adapted to live both on land and in water. This type of vernal pool is restricted to small, irregular depressions in tertiary pyroclastic flows. In this case, the restrictive layer is composed of volcanic mudflow material. Typical plant species encountered in these vernal pool communities include smooth goldfields (*Lasthenia glabberima*), bristled downingia (*Downingia bicornuta*), Douglas’ meadowfoam (*Limnanthes douglasii*), stalked popcornflower, white-headed navarretia (*Navarretia leucocephala*), pigmy-weed, and woolly marbles (*Psilocarpus brevissimus*). Northern volcanic mudflow vernal pool is a community in the MSCS habitat type “natural seasonal wetland.”

Pool size and the depth, duration, and seasonal timing of ponding are important factors that influence the composition and diversity of plant and animal species in vernal pools (Solomeshch et al. 2007). Consequently, the vegetation of vernal pools can vary substantially from year to year in response to interannual fluctuations in climate.

Management activities such as grazing and burning also influence species composition and diversity. In fact, recent research indicates that the abundance

of nonnative grasses, grazing practices, and hydrology are strongly interrelated and can substantially affect the plant communities of vernal pools (Robins and Vollmar 2002, Pyke 2004, Marty 2005).

Seasonal Wetlands Seasonal wetlands are ephemeral wetlands that pond or remain flooded for long periods during a portion of the year, generally the rainy winter season, then dry up, typically in spring. They often occur in shallow depressions on flood terraces that are occasionally to infrequently flooded. Seasonal wetlands are herbaceous communities typically characterized by species adapted for growth in both wet and dry conditions, and may contain considerable cover of upland species as well. Species commonly present in seasonal wetlands include tall flatsedge, dallisgrass (*Paspalum dilatatum*), Bermudagrass (*Cynodon dactylon*), Italian ryegrass, Mediterranean barley (*Hordeum marinum* ssp. *gussoneanum*), and curly dock (*Rumex crispus*).

Seasonal wetlands differ from vernal pools in that they do not have a restrictive hardpan layer and are usually dominated by nonnative plant species, especially nonnative grasses. Vernal pools are typically distinguished by a unique host of native and endemic plant species adapted to the extreme conditions created by the cycles of inundation and drying. Seasonal wetlands differ from freshwater marshes and seeps in that they are not permanently flooded or saturated. The seasonal wetland community type is not included in the Holland or Sawyer and Keeler-Wolf classification systems, but it is recognized by USACE and may be subject to USACE jurisdiction under Section 404 of the CWA. Seasonal wetland communities are in both the MSCS “natural seasonal wetland” and “managed seasonal wetland” habitat types.

Extended Study Area

The extended study area extends from RBDD south to the Delta and includes the lower Sacramento River, as well as the lower Feather and American Rivers downstream from SWP and CVP reservoirs. It also includes the Bay-Delta area and portions of the American River basin, San Joaquin River basin, and the water service areas of the CVP and SWP. (See Figure 1-4.)

This vast area includes a large number of plant communities. The types of plant communities include the following:

- Submerged aquatic vegetation
- Permanent wetlands (fresh, brackish, and saline; tidal and nontidal)
- Riparian scrubs, woodlands, and forests
- Vernal pools and other seasonal wetlands
- Grasslands
- Oak woodlands
- Montane hardwood and conifer forests

Shasta Lake Water Resources Investigation
Biological Resources Appendix – Botanical Resources and Wetlands Technical Report

- Coastal and desert scrubs
- Chaparrals




Figure 1-4. Central Valley Project and State Water Project Service Areas

A number of these natural plant communities occur in the primary study area and are described above. The other natural plant communities are described in the following sections, and in Mayer and Laudenslayer 1988; Sawyer and Keeler-Wolf 1995; and CALFED 2000b.

In addition to natural plant communities, plant communities of agricultural and urban areas occupy extensive areas in the extended study area. Agricultural plant communities include alfalfa and other irrigated pastures, field and row crops, orchards and vineyards, and rice. Plant communities in developed areas, as described in Mayer and Laudenslayer (1988), include those of golf courses and parks, ruderal (disturbed) areas, landscaped development, and small areas of natural plant communities. Currently, agricultural and urban vegetation occupies nearly 70 percent of the Central Valley, and more than 70 percent of terrestrial habitats in the Delta. This extensive conversion of natural vegetation to agricultural and urban vegetation has reduced the extent of some natural plant communities more than others.

Lower Sacramento River and Delta The roughly 300 miles of the Sacramento River can be subdivided into distinct reaches. These reaches are discussed separately below because the differences among them in topography, hydrology, and geomorphology strongly affect riparian vegetation and associated habitat functions. This section focuses on the reaches of the main stem Sacramento River from RBDD to Colusa, from Colusa to the Delta, and in the Delta. Each of these reaches is discussed individually along with the main tributaries and floodplain bypasses to the Sacramento River. (See the Fisheries and Aquatic Ecosystem technical report for more information.)

Sacramento River from RBDD to the Delta The Sacramento River from RBDD to the Delta is described below by reach from RBDD to Colusa, and Colusa to the Delta. Primary tributaries to the lower Sacramento River and floodplain bypasses are also described.

Red Bluff Diversion Dam to Colusa In this approximately 54-mile reach, the Sacramento River is classified as a meandering river, where relatively stable, straight sections alternate with more sinuous, dynamic sections (Resources Agency 2003). The active channel is fairly wide in some stretches and the river splits into multiple forks at many different locations, creating gravel islands often with riparian vegetation. Historic bends in the river are visible throughout this reach and appear as scars of the historic channel locations with the riparian corridor and oxbow lakes still present in many locations. Well-developed riparian woodland occurs in many locations. The channel remains active and has the potential to migrate in times of high water. Point bars, islands, high and low terraces, instream woody cover, early-successional riparian plant growth, and other evidence of river meander and erosion are common in this reach.

As is characteristic of this section of the lower Sacramento River, major physiographic features include floodplains, basins, terraces, active and remnant channels, and oxbow sloughs. These features, together with the historic and current hydrology and dynamic meander pattern of the Sacramento River, sustain a diverse array of riparian plant communities along the river channel, intermixed in a broad arable floodplain. Although most of the mature valley oak woodland and savanna and other mature riparian forest community types farther from the river's edge are now absent from much of the lower Sacramento River corridor, this meandering section of the Sacramento River nonetheless supports a variety of habitat types: annual grassland, blackberry scrub, Great Valley willow scrub, Great Valley cottonwood forest, Great Valley mixed riparian forest, Great Valley valley oak woodland, and freshwater marsh. All of these habitat types are described above under the "Upper Sacramento River (Shasta Dam to Red Bluff)."

Colusa to the Delta The general character of the Sacramento River changes quite drastically downstream from Colusa from a dynamic and active meandering channel to a confined, narrow channel restricted from migration. Surrounding agricultural lands encroach directly adjacent to the levees, which have cut the river off from most of its riparian corridor, especially on the eastern side of the river. Most of the levees in this reach are lined with riprap, allowing the river no erodible substrate and limiting the extent of riparian vegetation.

Urban and agricultural encroachment, development, dams, levees, and demand for fuel resulted in the removal and fragmentation of most riparian areas along the lower Sacramento River. Native perennial grasslands once covered vast areas in the region but have since been farmed or invaded by nonnative annuals. Low-lying areas in the region once were routinely flooded, replenishing nutrients and providing water to many portions of the region not situated along waterways. However, diking and construction of levees to protect agricultural lands and residential areas have changed this, and the extent of many vegetation communities that depend on regular floods has been greatly reduced. Vernal pools, important wetland resources that were historically abundant, have decreased dramatically with agriculture and development in the last two centuries.

Primary Tributaries to the Lower Sacramento River The primary tributaries to the lower Sacramento River are the American and Feather Rivers; each is described separately below.

Lower Feather River The aquatic ecosystem in the lower Feather River, down to the confluence with the Sacramento River at Verona, is influenced by the California Department of Water Resources' (DWR) Oroville Facilities. The upper extent is fairly confined by levees as the river flows through the city of Oroville. Downstream from Oroville, the Feather River is fairly active and meanders its way south to Marysville. However, this stretch is bordered by active farmland, which confines the river into an incised channel in certain

stretches and limits the width of riparian woodland. Some of this adjacent farmland is in the process of being restored to floodplain habitat with the relocation of levees to become setback levees.

Lower American River The lower American River (below Folsom and Nimbus Dams) is fairly low gradient. Most of the lower American River is surrounded by the American River Parkway, which preserves the surrounding riparian zone. The river channel does not migrate to a large degree because it has become deeply incised, leaving tall cliffs and bluffs adjacent to the river.

Sacramento River Floodplain Bypasses Multiple water diversion structures in the lower Sacramento River move floodwaters into floodplain bypass areas during high-flow events. These floodplain bypass areas – the Butte Basin, Sutter Bypass, and Yolo Bypass – provide broad, inundated floodplain habitat during wet years. Unlike other Sacramento River and Delta habitats, floodplains and floodplain bypasses are seasonally dewatered (as high flows recede).

Sacramento–San Joaquin River Delta The Delta comprises an area of approximately 750,000 acres divided into a number of islands by hundreds of miles of waterways. Surface-water resources in the Delta are influenced by the interaction of tributary inflows, tides, Delta hydrodynamics, and diversions and transfers. The Sacramento, San Joaquin, Mokelumne, Cosumnes, and Calaveras Rivers all discharge directly into the Delta.

Before reclamation, the Delta was inundated each year by winter and spring runoff, which changed channel geometry in response to flood conditions and tidal influence. Consequently, there were extensive areas of marsh in the Delta, and large areas of peat soils (Histosols) formed.

Nearly all of the Delta's marshland has since been reclaimed by agriculture, peat production, and urban and industrial uses. More than 1,000 miles of levees protect this reclaimed land (CALFED 2000c). Reclaiming the land exposed the previously submerged peat soils to the atmosphere, and they began to oxidize and diminish. As a result, reclaimed Delta islands have been subsiding, and some are now as much as 25 feet below sea level.

Agriculture dominates the Delta area, with agricultural lands occupying approximately 72 percent of the region's total land area (CALFED 2000c). Agricultural vegetation includes agricultural lands that are seasonally flooded and those that are not. Major crops and cover types in agricultural production include small grains (such as wheat and barley), field crops (such as corn, sorghum, and safflower), truck crops (such as tomatoes and sugar beets), forage crops (such as hay and alfalfa), pastures, orchards, and vineyards. The distribution of seasonal crops varies annually, depending on crop-rotation patterns and market forces. Recent agricultural trends in the Delta include an increase in the acreage of orchards and vineyards. General cropping practices

result in monotypic stands of vegetation for the growing season and bare ground in fall and winter. In areas not intensively cultivated, such as fallow fields, roads, ditches, and levee slopes, regular maintenance precludes the establishment of ruderal vegetation or native vegetation communities.

Grassland and ruderal (i.e., disturbed upland) communities are present throughout the Delta region. Historically, native grasslands and seasonal wetlands occurred in the Delta region, but vernal pools were not common. However, vernal pools that occur in grasslands along the margins of the Delta region, such as the Jepson Prairie Preserve, support a wide diversity of native plants. The grassland, ruderal, and seasonal wetland communities of the Delta are similar to those of the upper and lower Sacramento River portions of the study area.

Although about one-half of the Delta's historical wetland areas have been diked and drained, some small islands remain in a quasinatural state, as do some other areas with aquatic and wetland communities (e.g., "flooded islands" that were once reclaimed land, but have been abandoned after levee failures). The species composition and ecology of these riparian, marsh, and aquatic plant communities differ from the composition and ecology of communities in the upper and lower Sacramento River portions of the study area. These are all sensitive communities and are described below.

Riparian Communities Along the lower Sacramento River and in the Delta, riparian vegetation is characterized by narrow linear strips of trees and shrubs, in single- to multiple-story canopies. Tree canopies may be continuous or discontinuous, or absent altogether (as in riparian scrubs). These patches of riparian vegetation may be on or at the toe of levees (particularly in the Delta). Riparian communities in this region include cottonwood-willow woodland, Valley oak riparian woodland, riparian scrub, and willow scrub. These communities are described below.

Cottonwood-willow woodland Cottonwood-willow woodland typically can be found on channel islands, on levees, and along unmaintained channel banks of Delta sloughs and rivers. The riparian zone along leveed islands is usually very narrow, but more extensive riparian areas occur on in-channel islands or in other unleveed areas. Dominant understory species include black willow, sandbar willow, and shining willow. Other understory species include Himalayan blackberry, California blackberry, California button-willow, Indian hemp, California rose, coyote brush, and California black walnut. Herbaceous cover occurs where shrubs are sparse or absent and includes Santa Barbara sedge, hoary nettle, creeping wildrye, and numerous nonnative species. Cottonwood-willow woodland is a community in the MSCS NCCP habitat "valley/foothill riparian."

Valley oak riparian woodland Valley oak riparian woodland includes areas where the dominant overstory is valley oak. Associated species are similar

to those described for the cottonwood-willow woodland vegetation. Valley oak riparian woodland is a community in the MSCS NCCP habitat “valley/foothill riparian.”

Riparian scrub Riparian scrub is typically found on channel islands on levees and along narrow channel banks of creeks, waterways, and major tributaries in the Delta region. It is dominated by dense stands of shrubs, such as California button-willow, wild rose, Himalayan blackberry, and white alder. Where shrub cover is absent, herbaceous cover is often abundant and includes Indian hemp, yellow iris, centaury, vervain, umbrella sedge, creeping bent grass, bugleweed, and hedge-nettle. Riparian scrub also includes blackberry thickets, which intergrade with other riparian habitats. These thickets are characteristically monotypic stands of Himalayan blackberry, with scattered and isolated trees and shrubs, including coyote brush, sandbar willow, shining willow, and white alder. Blackberry thickets occur in association with ruderal habitats; however, an herbaceous understory is not evident within these thickets. Elderberry shrubs may also be associated with this community type. Riparian scrub is a community in the MSCS NCCP habitat “valley/foothill riparian.”

Willow scrub Willow scrub is a type of riparian scrub habitat dominated by willow species, particularly sandbar willow and young trees of other willow species, such as shining willow and black willow. In disturbed areas, willow scrub intergrades with blackberry vegetation. Willow scrub is a community in the MSCS NCCP habitat “valley/foothill riparian.”

In the Delta, riparian areas are frequently dominated by nonnative invasive species, particularly along levees and berms. The most abundant of these are giant reed (*Arundo donax*), blue gum eucalyptus (*Eucalyptus globulus*), fennel (*Foeniculum vulgare*), perennial pepperweed, black locust (*Robinia pseudoacacia*), and Himalayan blackberry. However, a number of other species, such as fig (*Ficus carica*) and tree-of-heaven (*Ailanthus altissima*), are locally problematic.

The dynamics of riparian communities along the lower Sacramento River and in the Delta are similar to those described for riparian communities along the upper Sacramento River. However, along the Sacramento River south of Colusa, in the flood bypasses, and in the Delta, the disturbances that remove riparian vegetation, or create newly exposed surfaces where riparian vegetation can establish, differ somewhat from those along the upper Sacramento River. In these downstream areas, disturbances related to meander migration are more limited, and anthropogenic (human-caused) disturbances, such as levee maintenance and trampling, are greater than those upstream. This is because of the close proximity to levees, extensive placement of bank protection, and greater human population.

Marsh Communities Tidal freshwater and brackish-water emergent marsh is dominated by clonal perennial plants, particularly bulrushes (tules,

Scirpus sp.), and to a lesser extent cattails (*Typha* sp.), common reed (*Phragmites australis*), and waterpepper (*Polygonum hydropiperoides*). Tules, cattails, and giant reed are emergent macrophytes, large (up to 10 feet in height) rhizomatous plants rooted in the substrate with stems (culms) above the water surface. This community occurs on instream islands and along mostly unvegetated, tidally influenced waterways. Tidal freshwater and brackish-water emergent marsh corresponds to the MSCS NCCP habitat “tidal freshwater emergent.”

In tidal marsh vegetation, vegetation structure and species richness are strongly influenced by disturbance (e.g., wave action, fire), the range of elevations present at a site, and salinity. Disturbance provides opportunities for annuals and short-lived perennials to regenerate and for additional species (also primarily clonal perennials) to colonize the site, and it creates structural diversity. Disturbances in Delta marshes result from erosion, human uses, and fires. Erosion of channel banks creates eroded “scallop” and slumped sections of bank that are colonized by a number of species (Hart and Hunter 2004). Human uses include the clearing of trails, construction and use of blinds, and maintenance of ponds and clearing of vegetation to create open water. The diversity of native plant species in such disturbed areas can be noticeably greater than in relatively undisturbed areas nearby. Fires occur in the marshes of some Delta islands (e.g., Lower Sherman Island). Overall, these fires reduce thatch and aboveground biomass, and increase the availability of nutrients; this benefits annual and short-lived perennial species and increases the diversity of marsh vegetation.

In the Delta’s tidal wetlands, the cover of woody species and the richness (i.e., number) of species increase with elevation, and dominance frequently shifts from California bulrush (*Scirpus californicus*) to softstem bulrush. (Softstem bulrush is deciduous and California bulrush is evergreen; thus, these two species probably differ in their hydraulic roughness during the winter, and in the habitat they provide during winter and spring.)

The species composition of tidal marshes changes at upper elevations from that found at middle elevations. In brackish marshes, upper elevations may support halophytes such as saltgrass (*Distichlis spicata*) and pickleweed (*Salicornia* species). A number of species typical of freshwater and saline marshes may grow together in these high marsh areas. Areas with higher soil salinities support pickleweed, saltgrass, fat-hen, and gumplant (*Grindelia humilis*). If salinity levels are lower in the high marsh areas, brass buttons (*Cotula coronopifolia*) and Baltic rush (*Juncus balticus*) may be more prevalent. Alternatively, emergent wetlands may intergrade with the woody vegetation of adjacent riparian areas. Most woody plants in this transitional zone are shrubs and vines, including red osier dogwood, buttonbush, and willows. In the northern and central Delta, a transition from mid-level marsh to woody vegetation occurs (e.g., at Delta Meadows State Park), and in the western Delta mid-level marsh may transition to either woody vegetation or to an upper elevation marsh dominated by halophytes.

At lower elevations, there is also a transitional zone between marsh and aquatic vegetation. In this zone there are fewer species of emergent plants, tule stems are at a lower density, and occasionally clumps of submerged aquatics exist. A characteristic feature of this transition zone is floating mats of plants that are rooted in the substrate but have creeping stems that are prostrate on the water. Native plants on the fringes of the marsh with this form of growth include creeping water primrose (both the native *Ludwigia peploides* ssp. *peploides* and the nonnative *L. p.* ssp. *montevidensis*), which has both native and nonnative subspecies, and floating pennywort (*Hydrocotyle ranunculoides*). This floating fringe may be absent, or discontinuous and narrow (less than 3 feet), or it may extend out across the water surface for 3–12 feet with plants rooted in the substrate at the marsh's edge, and floating as a mat over deeper water. Other, smaller species of pennywort (*Hydrocotyle umbellata*, *H. verticillata*) also grow at the marsh's edge, but tend to be on exposed muddy banks and flats.

Regional salinity gradients also affect the species composition of the Delta's tidal marshes. Delta waters generally have minimal salt concentrations, but the water is brackish in areas of the lower Delta (like Lower Sherman Island) that are close to Suisun Bay. Salinity levels in the soil are driven primarily by the interaction between the salinity concentration of tidal waters, local weather conditions, and the marsh vegetation itself (Atwater and Hedel 1976). The presence of a certain plant species within different marsh vegetation types is the result of individual physiological tolerances and competition between species. In general, larger monocots inhabit the lowest marsh surfaces, which are inundated by most high tides, and surfaces at or above high-tide levels are dominated by broadleaf species and a few species of smaller monocots. Marshes flooded by fresh or brackish water support a more diverse assemblage of species that generally tolerate low to moderate salinity concentrations (Atwater and Hedel 1976).

There is currently little cover of nonnative invasive species in Delta marshes. However, perennial pepperweed (*Lepidium latifolium*) has become abundant in the upper elevations of many marshes; it is extremely difficult to eradicate. Purple loosestrife (*Lythrum salicaria*) may be a problematic invader in the future; elsewhere in the Delta it has established patches in the tidal zone. Other species create localized problems: water hyacinth will root in tidal marsh, and black locust, giant reed, and Himalayan blackberry shade or encroach on the tidal zone at the base of levees and berms. Also, several nonnative species not generally considered invasive, such as dallisgrass (*Paspalum dilatatum*), are abundant within the tidal zone and may be reducing available habitat for native species.

Aquatic Communities Aquatic vegetation consists of submerged plants generally rooted in the substrate, whose stems may partially extend above the water surface (e.g., during flowering) and floating plants that are generally not rooted in the substrate. This section focuses on vascular plants because of their dominance of aquatic vegetation in much of the Delta.

Submerged plants generally die back to their stem bases, rhizomes, and/or other overwintering vegetative structures (e.g., turions) as water temperatures drop in the late fall. Throughout spring and summer, active growth increases stem biomass (i.e., standing crop) to a peak in early fall. Although these plants flower and produce seed, reproduction via vegetative propagules (e.g., turions, specialized buds, and stem fragments) is their primary means of reproduction. The availability of light (which decreases with depth), turbidity, and shade cast by overtopping vegetation can restrict submerged plants to relatively shallow areas. In the Delta (which has turbid waters), most submerged vegetation appears to be restricted to areas less than 5–10 feet deep. The velocity of flows may contribute to this depth restriction. Native species of submerged plants include coontail (*Ceratophyllum demersum*), common elodea (*Elodea canadensis*), waterbuttercup (*Ranunculus aquatilis*), and pondweeds (*Potamogeton* sp.). Nonnative species include curlyleaf pondweed (*Potamogeton crispus*), egeria (*Egeria densa*), parrotfeather (*Myriophyllum aquaticum*), and Eurasian milfoil (*Myriophyllum spicatum*).

Most floating plants also depend primarily on vegetative reproduction. The growth rate of most species, and consequently their abundance, increases in late spring and summer, and then diminishes in late fall to early spring. Species also produce overwintering buds, spores, and seeds. Native species of floating plants in the Delta include duckweeds (*Lemna* sp., *Spirodela* sp.) and mosquito ferns (*Azolla* sp.). In addition, the nonnative water hyacinth (*Eichhornia crassipes*) is widespread and abundant.

Aquatic plant communities are in the MSCS NCCP habitat “tidal perennial aquatic.”

San Joaquin River Basin to the Delta The San Joaquin River basin includes the Central Valley south of the Delta. It is drier than the Sacramento Valley, and flows into the Delta from the San Joaquin River are considerably lower than those from the Sacramento River. Inflows from the Merced, Tuolumne, and Stanislaus Rivers historically contribute more than 60 percent of the flows in the San Joaquin River. Flows in these river systems are highly altered and are managed for flood control and water supply purposes. Numerous dams, reservoirs, and diversions are located on these rivers and others in the San Joaquin system. Historically, approximately 70 percent of the river’s runoff has been diverted to the Friant-Kern and Madera Canals, primarily for agricultural uses.

The San Joaquin River region has many similarities to the Sacramento River region; however, the San Joaquin River region’s riparian regions are not and have never been as extensive as those found in the Sacramento River region, and a larger portion of the land in the San Joaquin River region is devoted to agriculture (CALFED 2000c). Riparian communities and wetlands have been reduced by water diversions, reclamation activities, and conversion to agricultural land uses (CALFED 2000c). Wetlands are situated in the northern

and western reaches in the region but are less abundant in other parts of the region. Almost 70 percent of the lowlands have been converted to irrigated agriculture, and remaining natural vegetation has been fragmented.

CVP/SWP Service Areas The Delta region provides water to most of California's agriculture and to urban and industrial communities across the state. Within the Delta, CVP and SWP pumping plants move water from the Delta to a system of canals and reservoirs for agriculture, municipal, industrial, and environmental uses in the Central Valley, the Bay Area, along the central coast, and portions of Southern California.

Agricultural and urban areas dominate much of these service areas. Agricultural plant communities include alfalfa and other irrigated pastures, field and row crops, orchards and vineyards, and rice. Plant communities in developed areas, as described in Mayer and Laudenslayer (1988), include those of golf courses and parks, ruderal (disturbed) areas, landscaped development, and small areas of natural plant communities.

Although agricultural and urban land uses have substantially reduced the area and connectivity of natural vegetation, the service areas still contain a large diversity of both lowland and upland plant communities, including many sensitive plant communities (CALFED 2000c, Attachment 1). The most dramatic difference between historical and existing conditions is the fragmentation of what were once large contiguous blocks of habitat, such as chamise-redshank chaparral, coastal sage scrub, grassland, oak woodland, oak savanna, southern oak woodland-forest, riparian woodland-forest, succulent scrub, sand dune habitat, alkali desert scrub, desert riparian habitat, desert wash, freshwater/saltwater marsh, and coastal strand (CALFED 2000c). Significant changes to the natural landscape in the region occurred in the late 1800s and early 1900s with land conversions to agriculture, a pattern similar to that in the Central Valley. However, that pattern shifted dramatically compared to the pattern in the Central Valley, as urban growth in the region that started in the 1900s began to convert large areas of agricultural lands and of remaining natural vegetation to developed land uses.

Special-Status Species

Special-status species addressed in this section include plants that are legally protected or are otherwise considered sensitive by Federal, State, or local resource conservation agencies and organizations. These include species that are State listed and/or Federally listed as rare, threatened, or endangered; those considered as candidates or proposed for listing as threatened or endangered; species identified by DFG as Species of Special Concern or USFS as sensitive, endemic, or needing additional survey or management actions; and plants considered jointly by DFG and CNPS to be rare, threatened, or endangered; and species afforded protection under local planning documents, including the CALFED MSCS.

Primary Study Area

Shasta Lake and Vicinity Within the Shasta Lake and vicinity portion of the primary study area are a wide variety of vegetative communities and habitat components that support a large diversity of plant species. To aid in determining the potential impacts of the project, a list of potential plant species of concern was developed.

For the purposes of this evaluation, botanical species of concern are plants, lichen, and fungi that fall into any of the following categories:

- Designated as rare or listed as threatened or endangered by the State or Federal government
- Proposed for designation as rare or listing as threatened or endangered by the State or Federal government
- Candidate species for State or Federal listing as threatened or endangered
- Ranked as California Rare Plant Rank (CRPR) 1B, 2, 3, or 4 (formerly CNPS List 1A, 1B, 2, 3, or 4)
- Considered sensitive or endemic by USFS
- Considered a survey and manage species by USFS or BLM
- Designated as an MSCS covered species by CALFED

Potentially occurring plant species of concern were determined by performing several database searches, reviewing USFWS and DFG special-status species lists for Shasta County, reviewing other appropriate literature, discussions with resource agency personnel, and professional experience in the region. Additionally, results from the various vegetation habitat mapping efforts, botanical surveys, and wildlife surveys conducted in the area by NSR since 2002 were used in developing the list of species of concern.

Table 1-3 summarizes special-status plant species identified as having a potential to occur in the Shasta Lake and vicinity portion of the primary study area.

Table 1-3. Plant Species of Concern with Potential to Occur in the Shasta Lake and Vicinity Portion of the Primary Study Area

Common Name	Scientific Name	Status ¹
Shasta ageratina	<i>Ageratina shastensis</i>	CRPR 1B.2, USFS E
Bent-flowered fiddleneck	<i>Amsinckia lunaris</i>	CRPR 1B.2, BLMS
Mallory's manzanita	<i>Arctostaphylos malloryi</i>	CRPR 4.3
Shasta County arnica	<i>Arnica venosa</i>	CRPR 4.2, USFS E
Depauperate milk-vetch	<i>Astragalus pauperculus</i>	CRPR 4.3
Moonwort, grape-fern	<i>Botrychium</i> subgenus <i>Botrychium</i>	USFS S, S&M
Yellow-twist horsehair	<i>Bryoria tortuosa</i>	BLMS
Green bug moss	<i>Buxbaumia viridis</i>	USFS S, BLMS, S&M
Callahan's mariposa lily	<i>Calochortus syntrophus</i>	CRPR 1B.1
Butte County morning-glory	<i>Calystegia atriplicifolia</i> ssp. <i>buttensis</i>	CRPR 4.2
Castle Crags harebell	<i>Campanula shetleri</i>	CRPR 1B.3, USFS S, BLMS
Buxbaum's sedge	<i>Carex buxbaumii</i>	CRPR 4.2
Bristly sedge	<i>Carex comosa</i>	CRPR 2.1, MSCS r
Fox sedge	<i>Carex vulpinoidea</i>	CRPR 2.2
Shasta clarkia	<i>Clarkia borealis</i> ssp. <i>arida</i>	CRPR 1B.1, MSCS m, BLMS
Northern clarkia	<i>Clarkia borealis</i> ssp. <i>borealis</i>	CRPR 1B.3, BLMS
Silky cryptantha	<i>Cryptantha crinita</i>	CRPR 1B.2, MSCS m, BLMS
California lady's-slipper	<i>Cypripedium californicum</i>	CRPR 4.2
Clustered lady's-slipper	<i>Cypripedium fasciculatum</i>	CRPR 4.2, USFS S, BLMS, S&M
Mountain lady's-slipper	<i>Cypripedium montanum</i>	CRPR 4.2, USFS S, BLMS, S&M
Four-angled spike rush	<i>Eleocharis quadrangulata</i>	CRPR, MSCS m
Butte County fritillary	<i>Fritillaria eastwoodiae</i>	CRPR 3.2, USFS S
Dubious pea	<i>Lathyrus sulphureus</i> var. <i>argillaceus</i>	CRPR 3
Broad-lobed linanthus	<i>Leptosiphon latisectus</i>	CRPR 4.3
Cantelow's lewisia	<i>Lewisia cantelovii</i>	CRPR 1B.2, USFS S, BLMS
Howell's lewisia	<i>Lewisia cotyledon</i> var. <i>howellii</i>	CRPR 3.2
Bellinger's meadowfoam	<i>Limnanthes floccosa</i> ssp. <i>bellingeriana</i>	CRPR 1B.2, MSCS m, BLMS
Awl-leaved navarretia	<i>Navarretia subuligera</i>	CRPR 4.3
Shasta snow-wreath	<i>Neviusia cliftonii</i>	CRPR 1B.2, USFS S, MSCS m, BLMS
Thread-leaved beardtongue	<i>Penstemon filiformis</i>	CRPR 1B.3, MSCS m, USFS S, BLMS
Narrow-petaled rein orchid	<i>Piperia leptopetala</i>	CRPR 4.3
Bidwell's knotweed	<i>Polygonum bidwelliae</i>	CRPR 4.3
Eel-grass pondweed	<i>Potamogeton zosteriformis</i>	CRPR 2.2, MSCS m
Pacific fuzzwort	<i>Ptilidium californicum</i>	USFS S, BLMS, S&M
Brownish beaked-rush	<i>Rhynchospora capitellata</i>	CRPR 2.2

Table 1-3. Plant Species of Concern with Potential to Occur in the Shasta Lake and Vicinity Portion of the Primary Study Area (contd.)

Common Name	Scientific Name	Status ¹
Sanford's arrowhead	<i>Sagittaria sanfordii</i>	CRPR 1B.2, MSCS m, BLMS
Marsh skullcap	<i>Scutellaria galericulata</i>	CRPR 2.2, MSCS m
Canyon Creek stonecrop	<i>Sedum obtusatum</i> ssp. <i>paradisum</i>	CRPR 1B.3, USFS S, BLMS
English Peak greenbriar	<i>Smilax jamesii</i>	CRPR 1B.3, USFS S, MSCS m, BLMS
Obtuse starwort	<i>Stellaria obtusa</i>	CRPR 4.3
Slender false lupine	<i>Thermopsis gracilis</i> var. <i>gracilis</i>	CRPR 4.3
Shasta huckleberry	<i>Vaccinium</i> sp., undescribed	Genetically distinct from coastal and Sierra Nevada populations; may warrant taxonomic consideration.
Oval-leaved viburnum	<i>Viburnum ellipticum</i>	CRPR 2.3

Notes:

CRPR = California Rare Plant Rank

USFS = U.S. Forest Service

¹Status Codes

CRPR 1B = Plants rare, threatened, or endangered in California and elsewhere

CRPR 2 = Plants rare, threatened, or endangered in California but more common elsewhere

CRPR 3 = Plants for which more information is need—a review list

CRPR 4 = Plants of limited distribution—a watch list

CRPR Threat Ranks

0.1 = Seriously threatened in California

0.2 = Fairly threatened in California

0.3 = Not very threatened in California

USFS E = USFS Endemic Species

USFS S = USFS Sensitive Species

S&M = Survey and Manage Species

MSCS = Multi Species Conservation Strategy covered species

R = Recovery. Recover species' populations within the MSCS focus area to levels that ensure the species' long-term survival in nature.

r = Contribute to recovery. Implement some of the actions deemed necessary to recover species' populations within the MSCS focus area.

m = Maintain. Ensure that any adverse effects on the species that could be associated with implementation of CALFED actions will be fully offset through implementation of actions beneficial to the species (CALFED 2000c).

BLMS = BLM sensitive

The CNDDDB was reviewed for records of special-status plant species in or near the Shasta Lake and vicinity portion of the primary study area. The CNDDDB is a database consisting of historical observations of special-status plant species, wildlife species, and natural communities. The CNDDDB is limited to reported sightings and is not a comprehensive list of special-status species that may occur in a particular area.

A search of the CNPS Electronic Inventory was also conducted. The Electronic Inventory allows users to query the database using a set of variable search criteria. The result of the search is a list of potentially occurring special-status plant species. The criteria used for the query included all CRPR 1A, 1B, 2, 3, and 4 plants (formerly CNPS) occurring in Shasta County in closed-cone

coniferous forest, chaparral, cismontane woodland, lower montane coniferous forest, marshes and swamps, pebble plain, valley and foothill grasslands, riparian forest, riparian woodland, and riparian scrub habitats between the elevations of approximately 900 feet and 2,500 feet.

Botanical Surveys Because botanical studies are ongoing, detailed technical memoranda describing methods, results, and conclusions will be provided in the FEIS.

NSR conducted several botanical surveys for special-status plant species in the Shasta Lake and vicinity portion of the primary study area. Botanical surveys were conducted in 2002, 2003, 2004, 2009, and 2010. A list of species observed during the surveys is provided as Attachment 2. Hickman (1993) was used as the standard reference for taxonomic nomenclature and identification. Special-status plant species detected during the surveys are discussed in more detail in Attachment 3.

The first botanical surveys were performed during 2002 along the Big Backbone and Squaw Creek arms. The surveys were conducted in accordance with the technical methods prescribed by Nelson (1994). In 2003, botanical surveys were conducted along 11 selected riverine reaches: Little Backbone Creek, Sugarloaf Creek, upper Sacramento River, middle Salt Creek, Salt Creek, Nosoni Creek, Dekkas Creek, Campbell Creek, Flat Creek, Ripgut Creek, and Potem Creek. The surveys were conducted in accordance with the technical methods prescribed by Nelson (1994). In 2004, botanical surveys were conducted at a series of randomly and nonrandomly selected locations. Nonrandomly selected sites were located throughout the Shasta Lake and vicinity portion of the primary study area (not including relocation areas) based on 2002 and 2003 survey results. Sites were selected based on the presence of unique habitat and ecological attributes, such as recently burned areas, unique geologic substrates, late-seral forests, and relatively rare plant series. Nonrandomly selected sites varied in size and often included several plant series types. Randomly selected sites were selected throughout the area using plant series polygons developed from previously completed vegetation mapping. Using geographic information systems (GIS), individual vegetation polygons were assigned a unique number, and 100 numbers (i.e., vegetation polygons) were then randomly selected.

Based on previous surveys resulting in discoveries of Shasta snow-wreath (*Neviusia cliftonii*) and Shasta huckleberry (*Vaccinium* sp. undescribed), specific surveys for these species were conducted in 2009 and 2010. These surveys were designed to identify potential habitat for and locate populations of these species outside of the proposed project area. Pedestrian surveys were conducted to search the focus areas identified. Using methods described in Lindstrand and Nelson (2006), potential survey areas were identified using soil and geologic information at known sites and choosing areas with those same characteristics. In addition, survey sites were identified using intuitive

techniques, such as selecting areas with vegetative cover types similar to those of known populations and areas near known populations (regardless of vegetative cover).

To address potential project impacts and evaluate potential mitigation measures, a genetic study of the Shasta snow-wreath was conducted in 2009 and 2010. The goal of the genetic study was to determine (1) whether all Shasta snow-wreath populations are genetically identical and (2) whether there are several homogeneous population clusters or whether some other pattern is present. Twenty-one of the 23 known Shasta snow-wreath sites were included in the study. The genetic study determined that the species is characterized by low genetic diversity and high levels of genetic differentiation (National Forest Genetics Laboratory 2010). No strong patterns were found between the Shasta snow-wreath populations and several physical and geographic variables, including soil, geology, population size, and geographic location. Although high levels of genetic differentiation and no strong population patterns are present, the genetic study found three general population clusters, providing insight and basic species information for potential mitigation planning.

Concurrently with the Shasta snow-wreath genetic study, a genetic study was conducted in 2009 and 2010 to determine whether the huckleberry is different genetically from red huckleberry and, if so, to determine if it warrants recognition as a new taxon. The genetic study determined that the species is genetically distinct from other huckleberry populations (National Forest Genetics Laboratory 2010). Based on the results of the genetic studies combined with phenotypic characteristics, this species warrants recognition as a new taxon. The Shasta huckleberry appears to be an uncommon and geographically restricted species.

In 2010, botanical surveys were conducted in all relocation areas including the dam footprint. The surveys were conducted in accordance with the technical methods prescribed by Nelson (1994).

Shasta County arnica (*Arnica venosa*), Northern clarkia (*Clarkia borealis* ssp. *borealis*), Butte County fritillary (*Fritillaria eastwoodiae*), Cantelow's lewisia (*Lewisia cantelovii*), Shasta snow-wreath, slender false lupine (*Thermopsis gracilis* var. *gracilis*), Shasta huckleberry, and oval-leaved viburnum (*Viburnum ellipticum*) were special-status plant species found both incidentally and during the surveys efforts.

One population of Shasta County arnica was found in ponderosa pine habitat south of Bridge Bay Resort along the Main Body and one population was found near the privately owned cabins on National Forest System lands in the Salt Creek inlet on the Sacramento Arm. Additionally, the USFS has located a population along the Sacramento Arm north of Slaughterhouse Island during surveys conducted in 2010 (Figure 1-5a and 1-5c).

One population of northern clarkia was found in hardwood-conifer/chaparral habitat near Bailey Cove on the McCloud Arm, and another population was found in hardwood-conifer/chaparral habitat in Sugarloaf Cove west of Beehive Point on the Sacramento Arm. Locations of northern clarkia found incidentally and during the surveys are shown in Figures 12-5c through 12-5d.

Surveys conducted by the USFS in 2010 located five populations of Butte County fritillary; two populations along Flat Creek and three populations along Ripgut Creek on the Pit Arm (Figure 1-5f).

One population of Cantelow's lewisia was discovered on a rock outcrop on the right bank of the upper Sacramento River near the Shasta Lake/upper Sacramento River transition zone. Additionally, the USFS has located three populations along Sacramento Arm near Elmore Mountain during surveys conducted in 2010 (Figure 1-5c).

Shasta snow-wreath is currently known from 23 locations, most of which occur at or near the periphery of Shasta Lake. Ten Shasta snow-wreath populations occur in habitats associated with limestone formations, and 13 occur in other habitat types. Most populations are associated with stream drainages or the lower portions of upland slopes. Of these, 13 Shasta snow-wreath populations were discovered during the botanical surveys along the McCloud Arm (south of Shasta Caverns and Keluche Creek), Pit Arm (Brock Creek, Ripgut Creek, Flat Creek, Stein Creek, and west of Stein Creek), and the Main Body (Blue Ridge east, Blue Ridge west, Blue Ridge middle, Cove Creek, south of Cove Creek, and Jones Valley). Locations of Shasta snow-wreath found incidentally and during the surveys are shown in Figures 1-5a through 1-5f.

Slender false lupine populations were discovered in all portions of the primary study area, generally on low-gradient slopes. Locations of slender false lupine found during the surveys and incidentally are shown in Figures 1-5a through 1-5f.

Shasta huckleberry is currently known from 12 locations in the upper Spring Creek, Dry Fork, (little) Squaw Creek, Shoemaker Gulch, and Little Backbone Creek drainages, and from the vicinity of Bully Hill. All locations occur in the area historically known as the Copper Belt of Shasta County in the immediate vicinity of historic copper mining activities. Shasta huckleberry occurs at four locations in the SLWRI project area: (little) Squaw Creek, Shoemaker Gulch, Little Backbone Creek, and Horse Creek near Bully Hill. Locations of Shasta huckleberry found incidentally and during the surveys are shown in Figures 1-5a through 1-5f.




Figure 1-5a. Special-Status Plant Species Occurring in Shasta Lake and Vicinity

This page left blank intentionally.




Figure 1-5b. Special-Status Plant Species Occurring in Shasta Lake and Vicinity

This page left blank intentionally.




Figure 1-5c. Special-Status Plant Species Occurring in Shasta Lake and Vicinity

This page left blank intentionally.




Figure 1-5d. Special-Status Plant Species Occurring in Shasta Lake and Vicinity

This page left blank intentionally.




Figure 1-5e. Special-Status Plant Species Occurring in Shasta Lake and Vicinity

This page left blank intentionally.




Figure 1-5f. Special-Status Plant Species Occurring in Shasta Lake and Vicinity

This page left blank intentionally.

Two oval-leaved viburnum populations were found during the surveys. One population was found in a forested upland slope west of Pine Point Campground along the McCloud Arm and a second population was found in chaparral habitat at Jones Valley along the Pit Arm near the Clikapudi Trail. Locations of oval-leaved viburnum found incidentally and during the surveys are shown in Figures 1-5d and 1-5f.

Extended Study Area

Most of the special-status plant species listed in Table 1-3 have the potential to occur within the extended study area. Numerous additional special-status plant species could occur in the extended study area. Attachment 4 of this technical report contains comprehensive lists of all sensitive plant species in the extended study area that have been reported to the CNDDB, or that otherwise have the potential to occur in the extended study area.

Lower Sacramento River and Delta A number of special-status plant species could be affected in the lower Sacramento River and Delta by changes in hydrology (CALFED 2000a). These include species associated with vernal pool, riparian, marsh, and aquatic plant communities; and several other species with restricted distributions on or near channel banks, active floodplains, flood bypasses, and Delta waterways. These assemblages of special-status species are described below.

Species of Vernal Pool Communities In addition to species that are potentially present in the primary study area (Table 1-3), special-status plant species that may be associated with vernal pools along the lower Sacramento River and in the Delta region include alkali milk-vetch (*Astragalus tener* var. *tener*), brittlescale (*Atriplex depressa*), Hoover's spurge (*Chamaesyce hooveri*), Contra Costa goldfields (*Lasthenia conjugens*), hairy orcutt grass (*Orcuttia pilosa*), slender Orcutt grass (*Orcuttia tenuis*), bearded popcornflower (*Plagiobothrys hystriculus*), Delta woolly-marbles (*Psilocarphus brevissimus* var. *multiflorus*), Crampton's tuctoria (*Tuctoria mucronata*), and Greene's tuctoria (*Tuctoria greenii*). The following is a description of each of these species:

- *Brittlescale* is a CNPS List 1B and MSCS "m" ("maintain") species. Suitable habitat for this annual plant consists of vernal pools, seeps, grassland, chenopod scrub, and playas. It is known from Alameda, Butte, Colusa, Contra Costa, Fresno, Glenn, Merced, Solano, Stanislaus, Tulare, and Yolo Counties. Threats to this species include habitat loss because of development, incompatible grazing practices, and trampling (CNPS 2011).
- *Alkali milk-vetch* is a CNPS List 1B and MSCS "r" ("contribute to recovery") species. Suitable habitat for this annual plant consists of moist, level settings on alkaline soils with high clay content. It has been observed in varied vegetation including grassland, borders of drainages, fallow rice fields, and among weedy grasses and forbs. Although once

more widely distributed, its current known distribution is centered in Solano County, and it also is known from Alameda, Merced, Napa, and Yolo counties. Several occurrences are on public land managed for game species. Threats to this species include habitat loss, incompatible grazing and game management practices (e.g., inundation of land for waterfowl during the growing season of Ferris's milkvetch), competition from the nonnative plants that now dominate much of the vegetation in which Ferris's milk vetch occurs, and small population sizes (10–400 individuals) that leave Ferris's milkvetch vulnerable to extirpation from chance events (CNDDDB 2007; CNPS 2011).

- *Hoover's spurge* is Federally listed as threatened and is a CNPS List 1B and MSCS "m" species. Suitable habitat for this annual plant consists of relatively large, deep vernal pools among the rolling hills, remnant alluvial fans, and depositional stream terraces at the base of the Sierra Nevada foothills. This species is now concentrated primarily in the Vina Plains in Tehama and Butte Counties, with other occurrences in Glenn County as well as the Visalia-Yettem area of Tulare County and the Hickman–La Grange area of Stanislaus County. Critical habitat for this species occurs in the extended study area in Stanislaus, Merced, and Tulare Counties. Threats to this species include habitat loss, incompatible agricultural and grazing practices, and nonnative species (CNDDDB 2007; CNPS 2011).
- *Contra Costa goldfields* is Federally listed as endangered and is a CNPS List 1B and MSCS "m" species. Suitable habitat for this annual plant consists of vernal pools and seasonally wet areas within woodland, alkaline playa, and valley and foothill grassland communities. The historic distribution of Contra Costa goldfields was once more extensive, but it is still known from at least one location in seven counties: Alameda, Contra Costa, Marin, Monterey, Napa, Solano, and Sonoma Counties. Threats to this species include habitat loss, hydrological alterations, incompatible grazing practices and nonnative plants (CNPS 2011).
- *Hairy orcutt grass* is Federally listed as endangered and is a CNPS List 1B and MSCS "m" species. Suitable habitat for this annual plant consists of vernal pools in rolling topography on remnant alluvial fans and stream terraces in the Central Valley. The historical range included the eastern margins of Sacramento and San Joaquin Valleys from Tehama County south to Stanislaus County and through Merced and Madera Counties. This species is now concentrated primarily in the Vina Plains in Tehama County, with an isolated occurrence in central Butte County and other occurrences in Glenn, Madera, and eastern Stanislaus Counties. Critical habitat for this species occurs in the extended study area in Tehama, Stanislaus, Merced, and Madera Counties. Threats to this species include habitat loss, incompatible

grazing practices, nonnative species, and trampling (CNDDDB 2007; CNPS 2011).

- *Slender orcutt grass* is Federally listed as threatened and is a CNPS List 1B and MSCS “m” species. Suitable habitat for this annual plant consists of vernal pools on remnant alluvial fans and high stream terraces and recent basalt flows in valley grassland and blue oak woodland. It has some ability to colonize artificial habitats, such as the margins of stock ponds. This species is now concentrated primarily in the vicinity of Dales, Tehama County. A secondary area of concentration is the Modoc Plateau in Lassen, Plumas, Shasta, and Siskiyou Counties, with a few occurrences in the Lake-Napa and Southeastern Sacramento Valley Vernal Pool Regions. Critical habitat for this species occurs in the extended study area in Tehama County. Threats to this species include habitat loss, incompatible grazing practices, nonnative species, and off-road vehicle use and trampling (CNDDDB 2007; CNPS 2011).
- *Bearded popcornflower* is a CNPS List 1B species. This species was rediscovered in 2005 after not being seen since 1892. Suitable habitat for this annual plant consists of vernal pools, vernal swales, and other moist areas in grassland. It is currently known from vernal pools and mesic grasslands in Solano County. Threats to this species include habitat loss, discing of remaining habitat, and competition from the nonnative plants that now dominate much of the vegetation in which bearded popcornflower occurs (CNPS 2011).
- *Delta woolly-marbles* is a CNPS List 4 species. Suitable habitat for this annual plant consists of vernal pools, vernal swales, and other moist areas in grassland. This vernal pool species is currently known from Alameda, Napa, San Diego, San Joaquin, Santa Clara, Solano, Stanislaus, and Yolo Counties (CNPS 2011).
- *Crampton’s tuctoria*, also known as Solano grass, is Federally listed and State listed as endangered, and is a CNPS List 1B and MSCS “r” species. Crampton’s tuctoria is an annual plant of vernal pools. This species is thought to have once grown in isolated parts of the northern Delta in areas that flooded during the wet season, but any former habitat there has been long since reclaimed for agriculture. Only a few individuals of the plant now exist, mostly in Yolo County. Threats to this species include habitat loss, incompatible grazing practices, and small population sizes (10–400 individuals) that leave Crampton’s tuctoria vulnerable to extirpation from chance events (CNDDDB 2007; CNPS 2011).
- *Greene’s tuctoria* is Federally listed as endangered and is a CNPS List 1B and MSCS “m” species. Suitable habitat for this annual plant

species consists of vernal pools in valley grasslands. This species is currently found in widely separated occurrences in Butte, Merced, Shasta, and Tehama Counties. Sixty percent of the extant occurrences are in the Vina Plains area of Tehama and Butte Counties. Eastern Merced County has about 30 percent of the known occurrences. Other occurrences are located in Glenn and Shasta Counties. The species has been extirpated from Fresno, Madera, San Joaquin, Stanislaus, and Tulare Counties. Critical habitat for this species occurs in the extended study area in Tehama, Stanislaus, Tuolumne, Merced, Mariposa, and Madera Counties. Threats to this species include habitat loss, and incompatible grazing and agricultural practices (CNDDDB 2007; CNPS 2011).

Species of Riparian and Marsh Communities In addition to species considered potentially present in the primary study area (Table 1-3), special-status plant species potentially associated with riparian and marsh communities along the lower Sacramento River and in the Delta region include bristly sedge (*Carex comosa*), Suisun thistle (*Cirsium hydrophilum* var. *hydrophilum*), Soft bird's-beak (*Cordylanthus mollis* ssp. *mollis*), Delta button-celery (*Eryngium racemosum*), Northern California black walnut (*Juglans hindsii*), Delta tule pea (*Lathyrus jepsonii* var. *jepsonii*), Mason's lilaeopsis (*Lilaeopsis masonii*), Delta mudwort (*Limosella subulata*), Sanford's arrowhead (*Sagittaria sanfordii*), Marsh skullcap (*Scutellaria galericulata*), blue skullcap (*Scutellaria lateriflora*), and Suisun Marsh aster (*Symphytotrichum lentum*). The following is a description of each of these species:

- *Bristly sedge* is a CNPS List 2 and MSCS “r” species. This perennial, rhizomatous herb grows on lake edges and wet places in marshes and swamps across much of North America, but it is most abundant in the glaciated regions of Eastern North America (Flora of North America Editorial Committee 2002). It is widely distributed but rarely collected in California; it is known from Contra Costa, Lake, Mendocino, Sacramento, Santa Cruz, Shasta, San Joaquin, and Sonoma Counties. The conversion of wetlands to other land uses may have contributed to the rarity of bristly sedge in the Delta. Threats to this species include additional habitat loss, marsh drainage, and other alterations to hydrology, as well as road maintenance activities (CNPS 2011).
- *Suisun thistle* is Federally listed as endangered and is a CNPS List 1B and MSCS “R” (“Recovery”) species. Habitat for this species is restricted to a narrow tidal band within large saltwater or brackish tidal marshes that have fully developed tidal channel networks. This species does not generally occur in fringe tidal marshes that are less than 300 feet wide (USFWS 1995). This species was thought to be extinct until it was rediscovered on Grizzly Island in 1989 (CNPS 2011). There are only two known occurrences of Suisun thistle remaining, both within Suisun Marsh in Solano County. Threats to this species include

alterations to marsh hydrology, nonnative species, and trampling (CNPS 2011).

- *Soft bird's-beak* is Federally listed as endangered and State listed as rare, and is a CNPS List 1B and MSCS "R" species. This hemiparasitic annual species grows in saline and brackish marshes and is restricted to a narrow band, typically in a marsh's higher elevations. The species is known from Contra Costa, Napa, and Solano Counties. The remaining populations range in size from a single individual to more than 100,000 plants. Within populations the number of individuals fluctuates considerably from year to year, often by more than an order of magnitude. Conversion of wetlands to other land uses has contributed to the decline of soft bird's-beak. Current threats to the remaining populations include competition from nonnative plants (in particular perennial pepperweed, *Lepidium latifolium*), alteration of wetland hydrology (including trenching of wetlands for mosquito abatement and diking), grazing and trampling, and erosion (USFWS 1995; CNDDDB 2007; CNPS 2011).
- *Delta button-celery* is a CNPS List 1B and MSCS "r" species. This biennial plant grows in open riparian scrub on river floodplains, primarily along the San Joaquin River; most occupied habitat receives periodic inundation. The species is known from Calaveras, Contra Costa, Merced, Stanislaus, and San Joaquin Counties. Threats to this species include conversion of habitat to agricultural land uses, flood control, incompatible grazing practices, and various anthropogenic disturbances (e.g., off-road vehicles, road maintenance) (CNDDDB 2007; CNPS 2011).
- *Northern California black walnut* is a CNPS List 1B and MSCS "r" species. Since the 1850s this tree species has experienced significant changes in its distribution, abundance, and the genetic composition of its populations. Before the 1850s native stands of Northern California black walnut appear to have been confined to three areas. The most extensive of these areas was along the lower Sacramento River in the vicinity of Walnut Grove (including Andrus and Grand Islands and adjacent islands). However, no pre-1850 individuals remain along the lower Sacramento River (Jepson 1910, Smith 1912). Despite practically being eliminated from an initially restricted distribution, Northern California black walnut has become widely distributed and abundant. Currently, Northern California black walnut occurs in riparian zones, on agricultural lands, along roadsides and as an urban "weed tree" throughout the Sacramento Valley and beyond (Griffin and Critchfield 1972; Resources Agency 2003). This change in its distribution was brought about by widespread planting as an ornamental, and its use as a rootstock for the English walnut (*Juglans regia*), and its subsequent dispersal and establishment within natural and ruderal vegetation. The

initial plantings of Eastern black walnut, and the subsequent and extensive cultivation of English walnut, has led to hybridization of Northern California black walnut with these species. Therefore, although most existing Northern California black walnut trees closely resemble the early descriptions and collections of this species, their ancestry to some extent is likely to include other walnut species (CNPS 2011).

- *Delta tule pea* is a CNPS List 1B and MSCS “r” species. This perennial vine typically grows above the tidal zone in riparian scrub or at the edges of riparian forest, although it also occurs in marsh vegetation. The species is known from Contra Costa, Napa, Sacramento, San Joaquin, Solano, and Sonoma Counties. The decline of Delta tule pea populations has resulted primarily from habitat loss, installation of riprap and maintenance of levees, hydrologic alterations, and competition from nonnative invasive plants. These continue to threaten the species (CNDDDB 2007; CNPS 2011).
- *Mason’s lilaeopsis* is a CNPS List 1B and MSCS “R” species. This diminutive (less than 10 inches), rhizomatous perennial grows within the tidal zone on open sites along shores and eroding banks, at the toes of cut banks, and in adjacent marshes (Mason 1957, CNDDDB 2007). The species is known from Alameda, Contra Costa, Marin, Napa, Sacramento, San Joaquin, and Solano Counties. The decline of Mason’s lilaeopsis populations has resulted primarily from habitat loss, altered hydrodynamics, installation of riprap and maintenance of levees, and accelerated erosion (resulting in part from boat wakes). These continue to be the primary threats affecting the species (CNDDDB 2007; CNPS 2011).
- *Delta mudwort* is a CNPS List 2 and MSCS “r” species. Like Mason’s lilaeopsis, this diminutive (less than 10 inches), rhizomatous perennial grows within the tidal zone on open sites along shores, on eroding banks, at the toes of cut banks, and in adjacent marshes (Mason 1957, CNDDDB 2007). The species is known from Contra Costa, Marin, Sacramento, San Joaquin, and Solano Counties. The decline of Delta mudwort populations has resulted primarily from habitat loss, altered hydrodynamics, installation of riprap and maintenance of levees, and accelerated erosion (resulting in part from boat wakes). These continue to be the primary threats affecting the species (CNDDDB 2007; CNPS 2011).
- *Sanford’s arrowhead* is a CNPS List 1B and MSCS “m” species. This rhizomatous, perennial plant grows as an emergent in freshwater (nontidal) marshes, ponds, ditches, and various other shallow, slow-moving freshwater habitats. It is widely distributed in the Central

Valley but rare. Threats to this species include habitat loss, channel maintenance, and hydrologic alterations (Hickman 1993; CNPS 2011).

- *Marsh skullcap* and *blue skullcap* are CNPS List 2 and MSCS “m” species. These rhizomatous perennial herbs are widely distributed in wetlands across North America, but are rare in California. These species are known from Sacramento and San Joaquin Counties in the Delta, and in California marsh skullcap also occurs in several counties in the northern Sierra Nevada and Modoc Plateau. Marsh skullcap is known from just three sites in the Delta, and all were growing in tidal marsh vegetation. Blue skullcap is known from just two sites. The conversion of wetlands to other land uses may have contributed to the rarity of these species in the Delta (CNDDDB 2007; CNPS 2011).
- *Suisun Marsh aster* is a CNPS List 1B species. This perennial grows primarily within the tidal zone, in patches of one to several hundred stems. However, although it does grow along banks, it also often is away from the water’s edge and within vegetation denser than vegetation surrounding Mason’s lilaeopsis or Delta mudwort. The decline of Suisun aster populations has been caused primarily by habitat loss, installation of riprap and maintenance of levees, hydrologic alterations, and competition from nonnative invasive plants. These continue to be the primary threats affecting the species (CNDDDB 2007; CNPS 2011).

Species of Aquatic Communities Eel-grass pondweed (*Potamogeton zosteriformis*) is a CNPS List 2 and MSCS “m” species. This submerged aquatic plant of assorted freshwater habitats is rare in California but more common elsewhere. In California, it has been documented in Contra Costa, Lake, Lassen, Modoc, and Shasta Counties. Overall, the distribution, abundance, and threats affecting this species in California are not well known. (CNPS 2011.)

CVP/SWP Service Areas Special-status plants are not likely to occur in a substantial portion of the CVP and SWP service areas because the agricultural and urban land uses tend to preclude suitable habitat for most native species.

General agricultural practices result in monotypic stands of vegetation for the growing season and bare ground in fall and winter. Special-status plant species are unlikely to occur in agricultural habitat because of the soil disturbance inherent in the agricultural practices. In areas not intensively cultivated, such as fallow fields, roads, ditches, and levee slopes, regular maintenance precludes the establishment of ruderal vegetation or native vegetation communities; thus, such areas are not likely to support many special-status plant species. Further, because agricultural ditches are generally subject to maintenance, they have minimal suitable habitat for special-status plants.

Developed land includes areas with roads and buildings but also includes barren areas that have been disturbed and are unvegetated. Special-status plant species are unlikely to occur in developed land areas because most vegetation has been removed, and these areas remain highly disturbed.

Areas with ruderal vegetation are dominated by herbaceous, nonnative, weedy species and may support stands of noxious weeds. Ruderal vegetation generally occurs in areas subject to periodic disturbances, and the species in this land cover type are generally weedy to invasive. For these reasons, no special-status plants are expected to occur in ruderal vegetation communities.

Although agricultural and developed land uses account for most of the CVP and SWP service areas, a portion of these areas still remains in natural vegetation. Because of the large size of the CVP and SWP service areas, this natural vegetation is distributed over a wide range of climate and soils, and is varied in structure and species composition. Consequently, a large number of special-status plant species has the potential to occur in the natural vegetation that remains within the CVP and SWP service areas (See Attachment 1).

Invasive Species

Primary Study Area

Shasta Lake and Vicinity Nonnative plant species introduced to the region by early settlers are of concern in the Shasta Lake and vicinity portion of the study area. When plants that evolved in one region of the globe are moved by humans to another region, a few flourish, crowding out native vegetation and wildlife that feed on the native species. Some invasive plants can even change ecosystem processes such as hydrology, fire regimes, and soil chemistry. These invasive plants have a competitive advantage because they are no longer controlled by their natural predators and can quickly spread. In California, approximately 3 percent of the plant species growing in the wild are considered invasive, but they inhabit a much greater proportion of the landscape (California Invasive Plant Council 2007).

Plant pests are defined by law, regulation, and technical organizations, and are regulated by many different bodies, including the California Department of Food and Agriculture (CDFA), U.S. Department of Agriculture (USDA), and the California Invasive Plant Council (CalIPC). CDFA uses an action-oriented pest-rating system. The low rating assigned to a pest by CDFA does not necessarily mean that the pest is not a problem; rather, the rating system is meant to prioritize response by CDFA and county agricultural commissioners. Plants on CDFA's highest priority "A" list are defined as plants "of known economic importance subject to state-county enforced action involving eradication, quarantine regulation, containment, rejection or other holding action."

CalIPC has developed a list of plant pests specific to California wildlands. The CalIPC list is based on information submitted by land managers, botanists, and researchers throughout the state and on published sources. To determine plant pests potentially occurring in the Shasta Lake and vicinity portion of the primary study area, this list was reviewed and local agencies (BLM, USFS, California Department of Transportation, and Shasta County Department of Agriculture) were contacted to gather knowledge of known weed locations (Table 1-4). Incidental observations of noxious weeds by NSR biologists and botanists were also recorded. Attachment 5 describes each weed source location, the potential mode of spread, and the risk of spread at each of the known sites.

Management actions have been required to prevent the loss of habitat caused by some of the more invasive exotic species that outcompete native vegetation. However, these management actions have been limited and have been confined primarily to areas adjacent to campgrounds and USFS facilities.

Table 1-4. Nonnative Plant Species Known to Occur in the Shasta Lake and Vicinity Portion of the Primary Study Area

Common Name	Scientific Name	CalIPC Rating ¹	CDFA Ranking ²	Habitat
Tree of heaven	<i>Ailanthus altissima</i>	Moderate	None	Grassland, oak woodland, riparian
Slender wild oats	<i>Avena barbata</i>	Moderate	None	Coastal scrub, grassland, oak woodland, forest
Common wild oats	<i>Avena fatua</i>	Moderate	None	Coastal scrub, grassland, oak woodland, forest
Rattlesnake grass	<i>Briza maxima</i>	Limited	None	Grassland
Ripgut brome	<i>Bromus diandrus</i>	Moderate	None	Dunes, scrub, grassland, woodland, forest
Soft brome	<i>Bromus hordeaceus</i>	Limited	None	Grassland, sage brush, serpentine soils
Red brome	<i>Bromus madritensis</i> ssp. <i>rubens</i>	High	None	Interior scrub, woodlands, grassland
Cheatgrass	<i>Bromus tectorum</i>	High	None	Interior scrub, woodlands, grassland
Lenspod whitetip	<i>Cardaria chalapensis</i>	Moderate-ALERT	B	Central Valley wetlands
Italian thistle	<i>Carduus pycnocephalus</i>	Moderate	None	Forest, scrub, grasslands, woodlands.
White knapweed	<i>Centaurea diffusa</i>	Moderate	A	Great Basin scrub, coastal prairie

**Table 1-4. Nonnative Plant Species Known to Occur in the Shasta Lake and Vicinity
 Portion of the Primary Study Area (contd.)**

Common Name	Scientific Name	CalIPC Rating ¹	CDFA Ranking ²	Habitat
Spotted knapweed	<i>Centaurea maculosa</i>	High	A	Riparian, grassland, wet meadows, forests
Yellow star-thistle	<i>Centaurea solstitialis</i>	High	C	Grassland, woodlands, occasionally riparian
Squarrose knapweed	<i>Centaurea virgata</i> var. <i>squarrosa</i>	Moderate	A	Scrub, grassland, pinyon-juniper woodland
Rush skeleton weed	<i>Chondrilla juncea</i>	Moderate	A	Grassland
Canada thistle	<i>Cirsium arvense</i>	Moderate	B	Grassland, riparian areas, forests
Bull thistle	<i>Cirsium vulgare</i>	Moderate	None	Riparian areas, marshes, meadows
Field bindweed	<i>Convolvulus arvensis</i>	Evaluated, not listed	C	Agricultural weed
Bermuda grass	<i>Cynodon dactylon</i>	Moderate	C	Riparian scrub, common landscape weed
Scotch broom	<i>Cystis scoparius</i>	High	C	Coastal scrub, oak woodland
Longbeak stork's bill	<i>Erodium botrys</i>	Evaluated, not listed	None	Many upland habitats
Redstem stork's bill	<i>Erodium cicutarium</i>	Limited	None	Many upland habitats
Leafy spurge	<i>Euphorbia esula</i>	High-ALERT	A	Forests, woodlands, juniper forests
Fig	<i>Ficus carica</i>	Moderate	None	Riparian woodland
Fennel	<i>Foeniculum vulgare</i>	High	None	Grassland, scrub
French broom	<i>Genista mospessulana</i>	High	C	Coastal scrub, oak woodland, grassland
English ivy	<i>Hedera helix</i>	High	None	Coastal forest, riparian areas
Mediterranean barley, foxtail	<i>Hordeum marinum</i> , <i>H. murinum</i>	Moderate	None	Grassland
Common St. John's wort	<i>Hypericum perforatum</i>	Moderate	C	Many habitats, disturbed
Dyer's woad, Marlahan mustard	<i>Isatis tinctoria</i>	Moderate	B	Great Basin scrub and grassland
Dalmation toadflax	<i>Linaria dalmatica</i>	Moderate	A	Grassland, forest clearings
Italian ryegrass	<i>Lolium multiflorum</i>	Moderate	None	Grassland, oak woodlands, pinyon-juniper woodland

Table 1-4. Nonnative Plant Species Known to Occur in the Shasta Lake and Vicinity Portion of the Primary Study Area (contd.)

Common Name	Scientific Name	CalIPC Rating ¹	CDFA Ranking ²	Habitat
Oleander	<i>Nerium oleander</i>	Evaluated, not listed	None	Riparian areas
Pokeweed	<i>Phytolacca americana</i>	None	None	Riparian forest, riparian woodland
Black locust	<i>Robinia pseudoacacia</i>	Limited	None	Riparian areas, canyons
Himalayan blackberry	<i>Rubus discolor</i>	High	None	Riparian areas, marshes, oak woodlands
Cutleaf blackberry	<i>Rubus laciniatus</i>	None	None	Riparian areas, marshes, oak woodlands
Curly dock	<i>Rumex crispus</i>	Limited	None	Grassland, vernal pools, meadows, riparian
Tansy ragwort	<i>Senecio jacobaea</i>	Limited	B	Grassland, riparian
Johnsongrass	<i>Sorghum halepense</i>	None	C	Disturbed sites, moist places
Spanish broom	<i>Spartium junceum</i>	High	None	Coastal scrub, grassland, wetlands, oak woodland, forests
Medusa-head	<i>Taeniatherum caput-medusae</i>	High	C	Grassland, scrub, woodland
Spreading hedgeparsley	<i>Torilis arvensis</i>	Moderate	None	Widespread
Common mullein	<i>Verbascum thapsus</i>	Limited	None	Meadows, riparian, sagebrush, pinyon-juniper woodland
Periwinkle	<i>Vinca major</i>	Moderate	None	Riparian, oak woodlands, coastal scrub
Rat-tail fescue	<i>Vulpia myuros</i>	Moderate	None	Coastal sage scrub, chaparral

Notes:

CalIPC = California Invasive Plant Council

CDFA = California Department of Food and Agriculture

¹ CalIPC Inventory Categories

High—Severe ecological impacts on physical processes, plant and animal communities, and vegetation structure. Reproductive biology and other attributes are conducive to moderate to high rates of dispersal and establishment. Widely distributed ecologically.

Moderate—Substantial and apparent ecological impacts on physical processes, plant and animal communities, and vegetation structure. Reproductive biology and other attributes are conducive to moderate to high rates of dispersal, although generally dependent on ecological disturbance. Ecological amplitude and distribution may range from limited to widespread.

Limited—Invasive but ecological impacts are minor. Reproductive biology and other attributes result in moderate rates of invasiveness. Ecological amplitude and distribution are generally limited, but may be locally persistent and problematic (California Invasive Plant Council 2007).

² CDFA Pest Ratings of Noxious Weed Species and Noxious Weed Seed

A – Eradication, containment, rejection, or other holding action at the state-county level.

B – Intensive control or eradication, where feasible, at the county level.

C – Control or eradication as local conditions warrant, at the county level.

Q – Rating as “A” is pending at the state or county level.

Upper Sacramento River (Shasta Dam to Red Bluff) and Lower Sacramento River and Delta A number of nonnative species have been introduced and become abundant in the riparian areas and marshes (fresh emergent wetlands) of the Sacramento Valley and Delta (Hunter et al. 2003). Several of these invasive nonnatives, including red sesbania (*Sesbania punicea*), Himalayan blackberry (*Rubus discolor*), giant reed (*Arundo donax*), and perennial pepperweed (*Lepidium latifolium*), form dense, monotypic stands that preclude the establishment of native species (Bossard et al. 2000). In general, these species displace native plants, reduce biodiversity, alter river flows, and reduce wildlife habitat values. Table 1-5 lists the most problematic of those species in Sacramento Valley and Delta riparian areas and marshes—invasive species rated “high” by CalIPC; these species have severe ecological impacts on physical processes, plant and animal communities, and vegetation structure (CalIPC 2006).

Table 1-5. Cal-IPC High-Rated Invasive Plants of Sacramento Valley and Delta Riparian and Marsh Habitats

Scientific Name Common Name	CalIPC Rating	Primary Riparian/Marsh Habitat(s)	Plant Type
<i>Arundo donax</i> Giant Reed	H	Riparian Forest/Scrub	Perennial Grass
<i>Cortaderia selloana</i> Pampas Grass	H	Riparian Scrub	Perennial Grass
<i>Foeniculum vulgare</i> Fennel	H	Riparian Scrub	Perennial Herb
<i>Lepidium latifolium</i> Perennial Pepperweed	H	Tidal and Nontidal Marsh, Riparian Scrub	Perennial Herb
<i>Lythrum salicaria</i> Purple Loosestrife	H	Tidal And Nontidal Marsh	Perennial Herb
<i>Rubus armeniacus</i> (= <i>R. discolor</i>) Himalayan Blackberry	H	Riparian Forest and Scrub, Nontidal Marsh	Vine
<i>Sesbania punicea</i> Red Sesbania	H, A	Riparian Forest and Scrub	Tree
<i>Tamarix chinensis</i> , <i>T. gallica</i> , <i>T. parviflora</i> , <i>T. ramosissima</i> Chinese Tamarisk, French Tamarisk, Small Flower Tamarisk, Salt Cedar	H	Riparian Forest and Scrub	Tree, Shrub

Source: Cal-IPC 2006

Cal-IPC Inventory Ratings

A = Alert – Plants with the potential to spread explosively; infestations currently small and localized.

H = High – These species have severe ecological impacts on physical processes, plant and animal communities, and vegetation structure. Their reproductive biology and other attributes are conducive to moderate to high rates of dispersal and establishment. Most are widely distributed ecologically.

Waters of the United States, including Wetlands, in Shasta Lake and Vicinity

NSR delineated waters of the United States (wetlands and other waters under Federal jurisdiction) in the impoundment area around the perimeter of Shasta Lake, and on public lands in the relocation areas. Cumulatively, 29,992 acres of Federal jurisdictional waters of the United States occur along the impoundment area, including Shasta Lake at full pool. Wetlands, totaling 25 acres, consist of fresh emergent/riparian wetland, intermittent swale, riparian wetland, seasonal wetland, seep/spring wetland, and vegetated ditch. Other waters, totaling 29,967 acres consist of ephemeral, intermittent, and perennial streams, roadside ditches, seep/spring other waters and lacustrine. Some demolition and construction activities associated with the impoundment area and relocation areas will extend into Shasta Lake below the existing full pool elevation. Therefore, the acreage

of the surface of Shasta Lake has been included with the acreage of waters of the United States for the impoundment area. Acreage totals for relocation areas will be provided in the FEIS.

The delineation was conducted in accordance with the routine on-site method identified in the *Corps of Engineers Wetlands Delineation Manual* (Environmental Laboratory 1987) (Corps Manual) and the *Interim Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Arid West Region* (USACE 2006) (Arid West Manual). Each on-site wetland determination was based on field observations of soil, vegetation, and hydrologic characteristics. Delineation of “other waters” was based on the presence of an ordinary high-water mark (OHWM) and whether the feature is tributary to waters of the United States. Data points were characterized and documented for 10 percent of all feature types along the perimeter of Shasta Lake. In each relocation area, at least one pair of data points was recorded for each feature type. Soil pits were dug to a depth sufficient to document the presence or confirm the absence of hydric soil or hydrology indicators. Indicator status of wetland plants was determined using the *National List of Plant Species That Occur in Wetlands: California Region 0* (Reed 1988). Positive indicators of hydric soils were observed in the field in accordance with the criteria outlined in *Field Indicators of Hydric Soils in the United States* (NRCS 2006). The hydric status of each soil map unit located in the study area was reviewed using the *Web Soil Survey* (Soil Survey Staff 2010). Indicators of depth and duration of soil saturation, ponding, and drainage patterns and the OHWM were observed in the field. The boundaries of each wetland feature and the three-parameter data points were mapped using rectified color aerial photography. A Trimble Pathfinder Pro XH Global Positioning System (GPS) capable of sub-foot accuracy was primarily used to delineate features in the relocation areas.

The fieldwork for the impoundment area was conducted in 2004, 2006, and 2010. Along the McCloud Arm, the impoundment area extends beyond the McCloud River Bridge. Fieldwork will be completed in this area during the spring/summer of 2011. The fieldwork in the relocation areas was completed in early January 2011 and is currently being processed. These data will be provided in the FEIS.

Main Body

The wetland delineation of the impoundment area along the Main Body was conducted from January to April 2010. Delineated waters of the United States consisted of wetlands (seep/spring and riparian wetland, and vegetated ditch) and other waters (ephemeral, intermittent, and perennial streams, seep/spring other waters, and roadside ditch). Table 1-6 specifies the acres of each of the types of jurisdictional waters that occur in the impoundment area.

Table 1-6. Jurisdictional Waters in the Impoundment Area

Jurisdictional Water Type	Area (Acres)					
	Main Body	Big Backbone Arm	Sacramento Arm	McCloud Arm	Squaw Creek Arm	Pit Arm
Wetlands						
Fresh emergent/riparian wetland	0.00	0.00	3.14	0.00	0.00	0.00
Intermittent swale	0.00	0.002	0.00	0.00	0.00	0.04
Riparian wetland	1.16	1.71	5.42	8.26	1.48	0.82
Seasonal wetland	0.00	0.00	0.18	0.00	0.14	0.02
Seep/spring wetland	0.77	0.23	0.80	0.31	0.13	0.41
Vegetated ditch	0.13	0.00	0.00	0.02	0.00	0.00
Total Wetlands	2.06	1.94	9.54	8.59	1.75	1.29
Other Waters of the United States						
Ephemeral stream	0.28	0.02	0.54	0.26	0.12	0.13
Intermittent stream	1.42	0.25	2.06	0.94	0.8	2.61
Perennial stream	1.53	3.00	8.67	20.27	2.29	1.46
Roadside ditch	0.00	0.00	0.02	0.00	0.00	0.00
Seep/spring other waters	0.03	0.00	0.001	0.01	0.00	0.00
Lacustrine	10,196.88	1,014.12	7,225.14	5,032.68	2,081.60	4,372.80
Total Other Waters	10,200.14	1,017.39	7,236.43	5,054.16	2,084.09	4,375.00
Total Waters of the U.S.	10,202.20	1,019.33	7,245.97	5,062.75	2,085.84	4,376.29

Note:

*Acreage values are approximate.

Big Backbone Arm

The wetland delineation along the Big Backbone Arm was conducted during November 2006. Delineated waters of the United States consisted of wetlands (seep/spring and riparian wetlands) and other waters (ephemeral, intermittent, and perennial streams). Table 1-6 specifies the acres of each of the types of jurisdictional waters occurring along the Big Backbone Arm.

Sacramento Arm

The wetland delineation along the Sacramento Arm was conducted primarily from September through early December 2010 and intermittently in March, April, and June 2010. Delineated waters of the United States consisted of wetlands (seep/spring, riparian, seasonal, and riparian/fresh emergent wetlands) and other waters (ephemeral, intermittent, and perennial streams, seep/spring

other waters, and roadside ditch). Table 1-6 specifies the acres of each of the types of jurisdictional waters occurring along the Sacramento Arm.

McCloud Arm

The wetland delineation along the McCloud Arm was conducted primarily during December 2009 and then intermittently in April, June, and November 2010. Delineated waters of the United States consisted of wetlands (seep/spring and riparian wetlands, and vegetated ditch) and other waters (ephemeral, intermittent, and perennial streams, and seep/spring other waters). Table 1-6 specifies the acres of each of the types of jurisdictional waters occurring along the McCloud Arm.

Squaw Creek Arm

The wetland delineation along the Squaw Creek Arm was conducted from late August through September 2004. Delineated waters of the United States include wetlands (seep/spring, riparian wetlands, and seasonal wet meadow) and other waters (ephemeral, intermittent, perennial streams, and seep/spring other waters). Table 1-6 specifies the acres of each of the types of jurisdictional waters occurring along the Squaw Creek Arm.

Pit Arm

The wetland delineation along the Pit Arm was conducted from late November 2006 through April 2007. Delineated waters of the United States consisted of wetlands (riparian, seep/spring, seasonal wetlands, and intermittent swale) and other waters (ephemeral, intermittent, and perennial streams). Table 1-6 specifies the acres of each of the types of jurisdictional waters occurring along the Pit Arm.

Characterization of Features

Wetlands mapped in the Shasta Lake and vicinity portion of the primary study area include fresh emergent/riparian wetland, intermittent swale, riparian wetland, seasonal wetland, seep/spring wetland, and vegetated ditch. The jurisdictional limit of each feature was delineated where all three wetland parameters (wetland vegetation, soils, and hydrology) were met.

One fresh emergent riparian wetland occurs along the Sacramento Arm at the confluence of Salt Creek and Shasta Lake. The I-5 crossing coupled with USFS attempts to develop the area for recreation has impounded the flows of Salt Creek, resulting in the development of fresh emergent riparian wetlands. Depending on water depths, wooded riparian areas and fresh emergent vegetation have established throughout the feature.

Dominant overstory species include Goodding's black willow (OBL¹), arroyo willow (FACW), red willow (assume FACW), and shining willow (OBL). Fresh

¹ OBL = Obligate Wetland Plants—Estimated probability of occurring in wetland >99 percent.
FACW = Facultative Wetland Plants—Estimated probability of occurring in wetland >67 percent to 99 percent.
FAC = Facultative Plants—Estimated probability of occurring in wetland 33 percent to 67 percent.

emergent species include pennyroyal (*Mentha pulegium*–OBL), willow dock (*Rumex salicifolius*–OBL), and broadleaf cattail (*Typha latifolia*). Inundation was observed during the field visit on December 16, 2010. Wetland hydrology and hydric soil criteria were met through evidence of frequent flooding, including sediment deposits, watermarks, drift lines, and drainage patterns.

Intermittent swales occur along the Big Backbone and Pit arms. Intermittent swales are characterized as linear, or somewhat linear, drainage features lacking evidence of scour, but where wetland plant species have established as a result of soil saturation. Typical species occurring in this feature include monkey flower (*Mimulus guttatus*–OBL), spiny fruit buttercup (*Ranunculus muricatus*–FACW), slender rush (*Juncus tenuis*–FACW), and centaury (*Centaureum venustum*–NL).

Riparian wetlands generally occur as “stringers,” or narrow features found only immediately adjacent to intermittent or perennial streams throughout the primary study area. Typical species found in riparian wetlands in the study area include arroyo willow (FACW), Goodding’s black willow (OBL), white alder (FACW), Oregon ash (FACW), Indian rhubarb (*Darmera peltata*–NL), mugwort (*Artemisia douglasiana*–FACW), California wild grape (FACW), and Himalayan blackberry (FACW). Wetland hydrology and hydric soil criteria are met through evidence of frequent flooding, including sediment deposits, watermarks, drift lines, and drainage patterns.

The seasonal wetlands occurring along the Sacramento, Squaw Creek, and Pit arms are influenced by adjacent water features or are depressions that frequently pond. Plant species found in these features include slender rush (FACW), sword leaf rush (*Juncus ensifolius*–FACW), monkey flower (OBL), yampah (*Perideridia californica*–FACW), annual checker bloom (*Sidalcea calycosa*–OBL), little quaking grass (*Briza minor*–FACW), California oatgrass (*Danthonia californica*–FACW), and spiny fruit buttercup (FACW). Wetland hydrology and soils criteria are met through evidence of long-duration saturation, including saturation in the upper 12 inches, aquic moisture regime, and drainage patterns.

Seep/spring wetlands are found throughout the primary study area; they form when groundwater flows out of the ground where the aquifer meets the ground surface. Hydrophytic vegetation colonizes the area where water is provided by the seep/spring. Typical species in these features include white alder (FACW), chain fern (*Woodwardia fimbriata*–FACW), goat’s beard (*Aruncus dioicus*–FACW), Indian rhubarb (NL), monkey flower (*Mimulus guttatus*–OBL), horsetail (*Equisetum arvense*–FAC), red stem dogwood (*Cornus stolonifera*–FACW), spicebush (NL), Himalayan blackberry (FACW), and western azalea

FACU = Facultative Upland Plants—Estimated probability of occurring in wetland 1 percent to <33 percent.

UPL = Obligate Upland Plants—Estimated probability of occurring in wetland <1 percent.

NI = No Indicator—Plants for which insufficient information was available to determine an indicator status.

NL = Not listed—Plants not listed in Reed 1988.

(FAC). The criteria for wetland hydrology and soils are met through evidence of long-duration saturation, including inundation, saturation in the upper 12 inches, watermarks, and drainage patterns.

There are few vegetated ditches in the Shasta Lake and vicinity area; they occur only along the Main Body and the McCloud Arm, and in relocation areas. Vegetated ditches are ditches that have been excavated to drain adjacent uplands, parking areas, roads, or railways. Because the gradients of these features are so slight, pooling and/or saturation occurs, allowing hydrophytic vegetation to colonize. Dominant plant species include nutsedge (*Cyperus eragrostis*–FACW), seep monkey flower (OBL), broadleaf cattail, and rush (*Juncus* sp.–assume FACW). Wetland hydrology criteria were met through the observation of surface water on March 19, 2010, and a thin muck surface indicating long-duration inundation. Hydric soil criteria were met through the observation of inundation and 1 centimeter (cm) of muck, indicating long-duration saturation. Other waters of the United States mapped in the Shasta Lake and vicinity portion of the primary study area include seep/spring other waters and ephemeral, intermittent, and perennial streams. The jurisdictional limit of each feature was delineated at the OHWM.

Ephemeral streams are characterized by indicators of scour and deposition, minor drift lines, and sediment deposits, but they lack a groundwater component that contributes to their flow. Hydrology is provided by sheet flow. The poorly defined hydrology indicators, proximity to the headwaters, and the small size of the ephemeral drainages indicate short duration of flow lacking a groundwater component.

Intermittent streams are the most abundant jurisdictional feature along the three arms. Intermittent streams range from small, poorly defined tributaries to larger, well-defined streams that flow into the summer. Like ephemeral streams, intermittent streams flow seasonally, but, in addition to precipitation and sheet flow from adjacent slopes, groundwater extends the duration of flow. Intermittent streams were identified as exhibiting the defined characteristics of a stream, generally a bed and bank and scour and depositions. Other characteristics, such as algae growth or hydrophytic vegetation in or adjacent to the stream, indicate inundation for a longer duration. Substrates observed were primarily cobble. Hydrology and hydric soil criteria are met through evidence of frequent flooding, including water marks, algal matting, drift lines, and sediment deposits.

Perennial streams exhibit the same characteristics as intermittent streams, but tend to be larger and to have a consistent source of groundwater. The substrate consists of boulders, bedrock, cobble, sand, and gravel. Riparian features often occur within the OHWM, but evident riparian “stringers” were delineated as wetlands. Hydrology and hydric soil criteria are met through evidence of frequent flooding, including water marks, algal matting, drift lines, and sediment deposits.

Roadside ditches are found along the Sacramento Arm near roadways and railroad tracks. These features have been excavated solely to drain uplands. Boundaries were established at the OHWM as indicated by sediment and drift deposits.

Seep/spring other waters are found along the Main Body, the Sacramento Arm, and the McCloud Arm. They form when groundwater flows out of the ground where the aquifer meets the ground surface. Hydrophytic vegetation is lacking. A channel may form, but in most cases water does not have the velocity to scour a bed and bank. Duff and organic soil substrates absorb the water and slow water movement downslope.

Regulatory Framework

Biological resources in California are protected and/or regulated by a variety of Federal and State laws and policies. In addition, in many parts of California, there are local or regional habitat and species conservation planning efforts in which a project applicant may participate. Key regulatory and conservation planning issues applicable to the project and alternatives under consideration are discussed below.

Federal

Endangered Species Act

Pursuant to the Federal Endangered Species Act (ESA), USFWS and the National Marine Fisheries Service (NMFS) have authority over projects that may result in “take” of a Federally listed species. In general, ESA Section 7 prohibits persons (including private parties) from “taking” listed endangered or threatened fish and wildlife species on private property, and from “taking” listed endangered or threatened plant species in areas under Federal jurisdiction or in violation of State law (16 U.S. Code (USC) 1532, 50 Code of Federal Regulations (CFR) 17.3). Under the ESA, the definition of “take” is to “harass, harm, pursue, hunt, shoot, wound, kill, trap, capture, or collect, or to attempt to engage in any such conduct” as part of an intentional or negligent act or omission. The term “harm” includes acts that result in death or injury to wildlife. Such acts may include significant habitat modification or degradation if it results in death or injury to wildlife by significantly impairing essential behavioral patterns, including breeding, feeding, or sheltering. Section 7(a) of the ESA, as amended, requires Federal agencies to evaluate their actions with respect to any species that is proposed for listing or is listed as endangered or threatened. Section 7(a)(2) requires Federal agencies to ensure that activities they authorize, fund, or carry out are not likely to jeopardize the continued existence of a listed species or to destroy or adversely modify its critical habitat. If a Federal action may affect a listed species or its critical habitat, the responsible Federal agency must enter into formal consultation with USFWS.

As defined in the ESA, critical habitat is a specific geographic area that is essential for the conservation of a threatened or endangered species and that may require special management and protection. It may include an area that is not currently occupied by the species but that will be needed for its recovery. Critical habitats are designated to ensure that actions authorized by Federal agencies will not destroy or adversely modify critical habitat, thereby protecting areas necessary for the conservation of the species.

Clean Water Act

The objective of the CWA is to restore and maintain the chemical, physical, and biological integrity of the nation's waters. USACE regulates discharges of fill or dredged materials into waters of the United States under Section 404 of the CWA (33 USC 1251–1376). Waters of the United States include lakes, rivers, streams, and their tributaries and adjacent wetlands. Wetlands are defined under Section 404 as areas that are inundated or saturated by surface water or groundwater at a frequency and duration sufficient to support (and that do support under normal circumstances) a prevalence of vegetation typically adapted for life in saturated soil conditions (33 CFR 328.3, 40 CFR 230.3). Activities that require a permit under Section 404 include, but are not limited to, placing fill or riprap, grading, mechanized land clearing, and dredging. Any activity that results in the deposit of dredged or fill material below the ordinary high-water mark of waters of the United States or within a jurisdictional wetland usually requires a Section 404 permit, even if the area is dry at the time the activity takes place. To comply with the Section 404 policy that there be no net loss of wetlands, the project cannot affect the total acreage of wetlands within the project boundary.

Section 401 Water Quality Certification

Under Section 401 of the CWA, an applicant for a Section 404 permit must obtain a certificate from the appropriate State agency stating that the intended dredging or filling activity is consistent with the State's water quality standards and criteria. In California, the authority to grant water quality certification is delegated by the State Water Resources Control Board to the nine regional water quality control boards (RWQCB).

Rivers and Harbors Act

USACE regulates the construction of structures in, over, or under; excavation of material from; or deposition of material into "navigable waters of the United States" under Section 10 of the Federal Rivers and Harbors Act (33 USC 401 et seq.). Navigable waters of the United States are defined as those waters subject to the ebb and flow of the tide shoreward to the mean high-water mark or those that are currently used, have been used in the past, or may be susceptible to use to transport interstate or foreign commerce. A letter of permission or permit from USACE is required before any work may be completed within navigable waters. Projects are permitted under either individual or general (i.e., nationwide) permits. The specific applicability of the permit types is determined by USACE on a case-by-case basis. Based on a preliminary conversation with

the USACE (San Francisco District, Eureka Field Office), the project is expected to be permitted under Nationwide Permit Number 27.

U.S. Forest Service Sensitive Species

The National Forest Management Act requires USFS to “provide for a diversity of plant and animal communities” (16 USC 1604(g)(3)(B)) as part of its multiple-use mandate. USFS must maintain “viable populations of existing native and desired nonnative species in the planning area” (36 CFR 219.19). The Sensitive Species program is designed to meet this mandate and to demonstrate USFS’s commitment to maintaining biodiversity on National Forest System lands. The program is a proactive approach to conserving species to prevent a trend toward listing under the ESA and to ensure the continued existence of viable, well-distributed populations. A “Sensitive Species” is any species of plant or animal that has been recognized by the Regional Forester to need special management to prevent the species from becoming threatened or endangered.

Shasta-Trinity National Forest Land and Resource Management Plan

The *Shasta-Trinity National Forest Land and Resource Management Plan* (STNF LRMP) contains forest goals, standards, and guidelines designed to guide the management of the STNF. The following goals, standards, and guidelines related to botanical resource issues associated with the study area were excerpted from the STNF LRMP (USFS 1995a).

U.S. Forest Service Survey and Manage

Standards and Guidelines In 1994, the U.S. Bureau of Land Management and USFS adopted standards and guidelines developed as part of the *Northwest Forest Plan*. These standards and guidelines address management of habitat for late-successional and old-growth forest related species within the range of the northern spotted owl. The *Northwest Forest Plan* was designed to address human and environmental needs served by the Federal forests of the western part of the Pacific Northwest and Northern California. The development of the *Northwest Forest Plan* was triggered in the early 1990s by the listing of the northern spotted owl and marbled murrelet as threatened under the ESA.

To mitigate potential impacts on plant and wildlife species that have the potential to occur within the range of the northern spotted owl, surveys are required for species thought to be rare, or whose status is unknown because of a lack of information. These species became known as the Survey and Manage species. The *Northwest Forest Plan* has gone through several revisions since its implementation in 1994, including the elimination of the Survey and Manage Mitigation Measure Standards and Guidelines in 2004. However, these guidelines were reinstated in January 2006 as the result of a court order.

Biological Diversity

Goals (LRMP, p. 4-4) Integrate multiple resource management on a landscape level to provide and maintain diversity and quality of habitats that support viable populations of plants, fish, and wildlife.

Standards and Guidelines (LRMP, p. 4-14)

- **Natural Openings** – Management of natural openings will be determined at the project level consistent with desired future conditions.
- **Snags** – Over time, provide the necessary number of replacement snags to meet density requirements as prescribed for each land allocation and/or management prescription. Live, green culls and trees exhibiting decadence and/or active wildlife use are preferred.
- **Hardwood** – Apply the following standards in existing hardwood types:
 - Manage hardwood types for sustainability.
 - Conversion to conifers will only take place to meet desired future ecosystem conditions.
 - Where hardwoods occur naturally within existing conifer types on suitable timber lands, manage for a desired future condition for hardwoods as identified during ecosystem analysis consistent with management prescription standards and guidelines. Retain groups of hardwoods over single trees.

Threatened, Endangered, and Sensitive Species (Plants and Animals)

Goals (LRMP, p. 4-5)

- Monitor and protect habitat for Federally listed threatened and endangered and candidate species. Assist in recovery efforts for threatened and endangered species. Cooperate with the State to meet objectives for state listed species.
- Manage habitat for sensitive plants and animals in a manner that will prevent any species from becoming a candidate for threatened and endangered status.

Botany (Sensitive and Endemic Plants)

Standards and Guidelines (LRMP, pp. 4-14 through 4-16)

- Map, record, and protect essential habitat for known and newly discovered sensitive and endemic plant species until conservation strategies are developed.

- Analyze the potential effects of all ground-disturbing projects on sensitive and endemic plants and their habitat. Mitigate project effects to avoid a decline in species viability at the Forest level.
- Monitor the effects of management activities on sensitive and endemic plants. If monitoring results show a decline in species viability, alter management strategy.
- Provide reports of sensitive plant populations to the DFG annually.
- Coordinate sensitive plant inventory and protection efforts with the DFG, the USFWS, the Nature Conservancy, the California Native Plant Society (CNPS), and other concerned agencies, organizations, and adjacent landowners.
- Protect type localities of sensitive and endemic plants for their scientific value.

Management Guide for the Shasta and Trinity Units of the Whiskeytown-Shasta-Trinity National Recreation Area

A portion of the Shasta Unit of the Whiskeytown-Shasta-Trinity National Recreation Area (NRA) is included in the Shasta Lake and vicinity portion of the primary study area. The Management Guide for the Whiskeytown-Shasta-Trinity NRA, including the Shasta Unit, contains management strategies intended to achieve or maintain a desired condition. These strategies take into account opportunities, management recommendations for specific projects, and mitigation measures needed to achieve specific goals. The following strategies related to botanical and wetland resource issues associated with the Shasta Lake and vicinity portion of the primary study area were excerpted from the Management Guide (USFS 1996).

Vegetation (Management Guide, pp. IV-18 through IV-19)

- Prescribed burning, fuel break construction, and other forms of vegetation manipulation will be used to reduce fire hazards and improve forest health.
- Hazard trees in traditionally high use recreation areas which pose safety hazards to people or property will be identified and removed.
- Recreation sites will be inventoried and vegetative management plans will be developed to ensure healthy and safe vegetation complexes are maintained over time.
- Protect known populations of Threatened and Endangered Species (TES) plant species and their habitat and implement mitigation measures if necessary to maintain or enhance their continued viability.

Conservation strategies for TES plant species will be utilized as they are developed.

- Implement management practices which promote restoration of native plant diversity.
- Implement a program to restore native vegetation to highly disturbed or degraded areas using native plants. Local in-kind, on-site seed or other propagation sources will be used in order to maintain genetic integrity.
- Chaparral and woodland habitat management will occur to meet wildlife objectives.
- Interpretive materials will address the need to conserve rare plant communities in accordance with the NRA Interpretive Plan.
- Rare plants in or near camping areas will be monitored on a regular basis.
- Diversity of native species will be emphasized. Eradication program will be implemented for nonnative, introduced species in areas where healthy, botanically diverse plant communities are necessary to meet ecosystem management objectives.
- Native plants from local gene pools will be utilized when landscaping campgrounds, interpretive facilities, etc.
- Partnerships will be utilized to assist with collection of seed, propagation of seeds/propagules, and planting.

U.S. Forest Service Noxious Weed Management Policy 2081

USFS Manual Policy 2080, Noxious Weed Management (USFS 1995b), includes a policy statement requiring a risk assessment for noxious weeds for every project. Specifically, the manual states:

2081.03. When any ground disturbing action or activity is proposed, determine the risk of introducing or spreading noxious weeds associated with the proposed action.

- *For projects having moderate to high risk of introducing or spreading noxious weeds, the project decision document must identify noxious weed control measures that must be undertaken during project implementation.*
- *Use contract and permit clauses to prevent the introduction or spread of noxious weeds by contractors and permittees. For example, where determined to be*

appropriate, use clauses requiring contractors or permittees to clean their equipment prior to entering National Forest System lands.

2081.2. Determine the factors which favor the establishment and spread of noxious weeds and design management practices or prescriptions to reduce the risk of infestation or spread of noxious weeds.

Where funds and other resources do not permit undertaking all desired measures, address and schedule noxious weed prevention and control in the following order:

- *First Priority: Prevent the introduction of new invaders,*
- *Second Priority: Conduct early treatment of new infestations, and*
- *Third Priority: Contain and control established infestations.*

Executive Order 11990: Protection of Wetlands

Executive Order 11990 established the protection of wetlands and riparian systems as the official policy of the Federal government. It requires all Federal agencies to consider wetland protection as an important part of their policies and take action to minimize the destruction, loss, or degradation of wetlands, and to preserve and enhance the natural and beneficial values of wetlands.

Executive Order 11312: Invasive Species

Executive Order 11312 directs all Federal agencies to prevent and control introductions of invasive nonnative species in a cost-effective and environmentally sound manner to minimize their economic, ecological, and human health impacts. Executive Order 11312 established a national Invasive Species Council made up of Federal agencies and departments and a supporting Invasive Species Advisory Committee composed of State, local, and private entities. The Invasive Species Council and Advisory Committee oversee and facilitate implementation of the Executive Order, including preparation of a National Invasive Species Management Plan.

State

California Endangered Species Act

Under the California Endangered Species Act (CESA), DFG has the responsibility for maintaining a list of endangered and threatened species (California Fish and Game Code, Section 2070). DFG also maintains a list of “candidate species,” which are species for which DFG has issued a formal notice that they are under review for addition to the list of endangered or

threatened species. In addition, DFG maintains lists of “species of special concern,” which serve as species “watch lists.” Pursuant to the requirements of CESA, an agency reviewing a proposed project within its jurisdiction must determine whether any State-listed endangered or threatened species may be present in the project study area and, if so, whether the proposed project would have a potentially significant impact on any of these species. In addition, DFG encourages informal consultation on any proposed project that may affect a species that is a candidate for state listing.

Project-related impacts on species listed as endangered or threatened under the CESA would be considered significant. State-listed species are fully protected under the mandates of the CESA. “Take” of protected species incidental to otherwise lawful management activities may be authorized under Section 2081 of the California Fish and Game Code. Under the CESA, “take” is defined as an activity that would directly or indirectly kill an individual of a species, but the definition does not include “harm” or “harass,” as the Federal act does. As a result, the threshold for take under the CESA is higher than that under the ESA.

Authorization from DFG would be in the form of an incidental take permit or as a consistency determination (Section 2080.1[a] of the Fish and Game Code). Section 2080.1[a] of the Fish and Game Code authorizes DFG to accept a Federal biological opinion as the take authorization for a state-listed species when a species is listed under both the ESA and the CESA.

California Native Plant Protection Act

The Native Plant Protection Act (California Fish and Game Code Sections 1900–1913) prohibits the taking, possessing, or sale within the state of any plants with a State designation of rare, threatened, or endangered, as defined by DFG. The NPPA’s definition of “endangered” and “rare” closely parallel the CESA definitions of “endangered” and “threatened” plant species. Project impacts on these species are not considered significant unless the species are known to have a high potential to occur within the area of disturbance associated with construction of the project.

Section 1602 of the California Fish and Game Code—Streambed Alteration

Diversions, obstructions, or changes to the natural flow or bed, channel, or bank of any river, stream, or lake in California that supports wildlife resources are subject to regulation by DFG, pursuant to Section 1602 of the California Fish and Game Code. The regulatory definition of stream is a body of water that flows at least periodically or intermittently through a bed or channel that has banks and supports wildlife, fish, or other aquatic life. This includes watercourses that have a surface or subsurface flow that supports or has supported riparian vegetation. DFG’s jurisdiction within altered or artificial waterways is based on the value of those waterways to fish and wildlife. A DFG streambed alteration agreement must be obtained for a project that would result in an impact on a river, stream, or lake.

Porter-Cologne Water Quality Control Act

The Porter-Cologne Water Quality Control Act requires that each of the nine RWQCBs prepare and periodically update basin plans for water quality control. Each basin plan sets forth water quality standards for surface water and groundwater and actions to control nonpoint and point sources of pollution to achieve and maintain these standards. Basin plans offer an opportunity to protect wetlands through the establishment of water quality objectives. The RWQCB's jurisdiction includes Federally protected waters as well as areas that meet the definition of "waters of the state." Waters of the state is defined as any surface water or groundwater, including saline waters, within the boundaries of the state. The RWQCB has the discretion to take jurisdiction over areas not Federally protected under Section 401 provided they meet the definition of waters of the state. Mitigation requiring no net loss of wetlands functions and values of waters of the state is typically required by the RWQCB.

California Department of Fish and Game Species Designations

DFG maintains an informal list of species called "species of special concern." These are broadly defined as plant and wildlife species that are of concern to DFG because of population declines and restricted distributions, and/or because they are associated with habitats that are declining in California. These species are inventoried in the CNDDDB regardless of their legal status. Impacts on species of special concern may be considered significant.

California Native Plant Society Species Designations

CNPS is a statewide nonprofit organization that seeks to increase understanding of California's native flora and to preserve this rich resource for future generations. DFG and CNPS assign rare plant ranks through the collaborative efforts of the Rare Plant Status Review Group composed of over 300 botanical experts from government, academia, non-government organizations, and the private sector and managed jointly by DFG and CNPS. California native plants meeting the rarity or endangerment criteria are assigned a California Rare Plant Rank (CRPR). These plants were formerly referred to as CNPS listed species; however, as of March 2010, DFG has adopted the name CRPR for the rarity and endangerment categories to eliminate the false impression that these assignments are the exclusive work of CNPS and that CNPS has had undue influence over the regulatory process. CRPR 1 and 2 species generally qualify as endangered, rare, or threatened within the definition of State CEQA Guidelines CCR Section 15380. In general, CRPR 3 and 4 species do not meet the definition of endangered, rare, or threatened pursuant to CEQA Section 15380; however, these species may be evaluated by the lead agency on a case-by-case basis to determine significance criteria under CEQA.

Local

Shasta, Tehama, Glenn, Sutter, Sacramento, and Yolo counties and the Cities of Redding, Colusa, and Sacramento have established codes and policies that address protection of natural resources, including vegetation, sensitive species, and trees, and are applicable to the project.

Shasta County's general plan emphasizes that the maintenance and enhancement of quality fish and wildlife habitat is critical to the recreation and tourism industry, and acknowledges that any adverse and prolonged decline of these resources could result in negative impacts on an otherwise vibrant industry. The general plan identifies efforts to protect and restore these habitats to sustain the long-term viability of the tourism and recreation industry (Shasta County 2004).

The City of Redding's general plan strives to strike a balance between development and conservation by implementing several measures such as creek-corridor protection, sensitive hillside development, habitat protection, and protection of prominent ridge lines that provide a backdrop to the city (City of Redding 2000).

Tehama County's general plan update provides an overarching guide to future development and establishes goals, policies, and implementation measures designed to address potential changes in county land use and development. The general plan identifies the importance of retaining agriculture as one of the primary uses of land in Tehama County.

Glenn County's general plan provides a comprehensive plan for growth and development in Glenn County for the next 20 years (2007–2027). This plan recognizes that public lands purchased for wildlife preservation generate economic activity as scientists and members of the public come to view and study remnant ecosystems (Glenn County 1993).

The City of Colusa's general plan seeks to promote its natural resources through increased awareness and improved public access (City of Colusa 2007).

Sutter County's general plan contains policies that generally address preservation of natural vegetation, including wetlands. It requires that new development mitigate the loss of Federally protected wetlands to achieve "no net loss," but it does not include any other specific requirements.

Sacramento County's general plan contains policies that promote protection of marsh and riparian areas, including specification of setbacks and "no net loss" of riparian woodland or marsh acreage (Sacramento County 1993). It also addresses the need to conserve vernal pools and ephemeral wetlands to ensure no net loss of vernal pool acreage. Several policies specifically promote protection of native oak trees, and, in some areas of the county, seek to ensure that there is no net loss of canopy area. The general plan for the County of Sacramento is currently under revision.

The City of Sacramento Municipal Code addresses the protection of trees within the city boundaries, including general protection of all trees on city property and specific protection of heritage trees.

Yolo County's general plan aims to provide an active and productive buffer of farmland and open space separating the Bay Area from Sacramento, and integrating green spaces into its communities.

Federal, State, and Local Programs and Projects

California Bay-Delta Authority

The California Bay-Delta Authority was established as a State agency in 2003 to oversee implementation of CALFED for the 25 Federal and State agencies working cooperatively to improve the quality and reliability of California's water supplies while restoring the Bay-Delta ecosystem. The CALFED Ecosystem Restoration Program has provided a funding source for projects that include those involving acquisition of lands within the Sacramento River Conservation Area, initial baseline monitoring and preliminary restoration planning, and preparation of long-term habitat restoration management and monitoring plans.

Cantara Trustee Council

The Cantara Trustee Council administers a grant program that has provided funding for numerous environmental restoration projects in the primary study area, including programs in the Fall River watershed, Sulphur Creek, the upper Sacramento River, Middle Creek, lower Clear Creek, Battle Creek, Salt Creek, and Olney Creek. The Cantara Trustee Council is a potential local sponsor for future restoration actions in the primary study area. The Cantara Trustee Council includes representatives from DFG, USFWS, the Central Valley Regional Water Quality Control Board, California Sportfishing Protection Alliance, and Shasta Cascade Wonderland Association.

Resource Conservation Districts

There are numerous resource conservation districts (RCD) within the study area. Once known as soil conservation districts, RCDs were established under California law with a primary purpose to implement local conservation measures. Although RCDs are locally governed agencies with locally appointed, independent boards of directors, they often have close ties to county agencies and the National Resources Conservation Service. RCDs are empowered to conserve resources within their districts by implementing projects on public and private lands and to educate landowners and the public about resource conservation. They are often involved in the formation and coordination of watershed working groups and other conservation alliances. In the Shasta Lake and upper Sacramento River vicinity, districts include the Western Shasta County RCD and the Tehama County RCD. To the east are the Fall River and Pit River RCDs, and to the west and north are the Trinity County and Shasta Valley RCDs.

Riparian Habitat Joint Venture

The Riparian Habitat Joint Venture (RHJV) was initiated in 1994 and includes signatories from 18 Federal, State, and private agencies. The RHJV promotes

conservation and the restoration of riparian habitat to support native bird population through three goals:

- Promote an understanding of the issues affecting riparian habitat through data collection and analysis.
- Double riparian habitat in California by funding and promoting on-the-ground conservation projects.
- Guide land managers and organizations to prioritize conservation actions.

RHJV conservation and action plans are documented in the *Riparian Bird Conservation Plan* (RHJV 2004). The conservation plan targets 14 “indicator” species of riparian-associated birds and provides recommendations for habitat protection, restoration, management, monitoring, and policy. The report notes habitat loss and degradation as one of the most important factors causing the decline of riparian birds in California. The RHJV has participated in monitoring efforts within the Sacramento National Wildlife Refuge Complex and other conservation areas. The RHJV’s conservation plan identifies lower Clear Creek as a prime breeding area for yellow warblers and song sparrows, advocating a continuous riparian corridor along lower Clear Creek. Other recommendations of the conservation plan apply to the North Delta Offstream Storage Investigation study area in general.

Sacramento River Advisory Council

In 1986 the California Legislature passed Senate Bill (SB) 1086, which called for a management plan for the Sacramento River and its tributaries to protect, restore, and enhance fisheries and riparian habitat in an area stretching from the confluence of the Sacramento River with the Feather River and continuing northward to Keswick Dam, about 4 miles north of Redding. The law established an advisory council that included representatives of Federal and State agencies, county supervisors, and representatives of landowners, water contractors, commercial and sport fisheries, and general wildlife and conservation interests. Responsibilities of the advisory council included development of the *Sacramento River Conservation Area Forum Handbook* to guide management of riparian habitat and agricultural uses along the river (Resources Agency 2003). This action also resulted in formation in May 2000 of the Sacramento River Conservation Area (SRCA) Forum, a nonprofit, public benefit corporation with a board of directors that includes private landowners and public interest representatives from a seven-county area, an appointee of the California Resources Agency, and ex-officio members from six Federal and State resource agencies. The work of the organization is generally focused on planning actions and river management within the SRCA planning area.

Sacramento River Conservation Area Program

SB 1086 called for a management plan for the Sacramento River and its tributaries to protect, restore, and enhance both fisheries and riparian habitat. The SRCA Program has an overall goal of preserving remaining riparian habitat and reestablishing a continuous riparian ecosystem along the Sacramento River between Redding and Chico, and reestablishing riparian vegetation along the river from Chico to Verona. The program is to be accomplished through an incentive-based, voluntary river management plan. The *Upper Sacramento River Fisheries and Riparian Habitat Management Plan*, January 1989 (Resources Agency 1989), identifies specific actions to help restore the Sacramento River fishery and riparian habitat between the Feather River and Keswick Dam. The *Sacramento River Conservation Area Forum Handbook* (Resources Agency 2003) is a guide to implementing the program. The Keswick Dam-to-Red Bluff portion of the conservation area includes areas within the 100-year floodplain, existing riparian bottomlands, and areas of contiguous valley oak woodland, totaling approximately 22,000 acres. The 1989 fisheries restoration plan recommended several actions specific to the study area:

- Fish passage improvements at RBDD (partially complete)
- Modification of the Spring Creek Tunnel intake for temperature control (completed)
- Spawning gravel replacement program (ongoing)
- Development of side-channel spawning areas, such as those at Turtle Bay in Redding (ongoing)
- Structural modifications to the Anderson-Cottonwood Irrigation District Dam to eliminate short-term flow fluctuations (completed)
- Maintaining instream flows through coordinated operation of water facilities (ongoing)
- Improvements at the Coleman National Fish Hatchery (partially complete)
- Measures to reduce acute toxicity caused by acid mine drainage and heavy metals (ongoing)
- Various fisheries improvements on Clear Creek (partially complete)
- Flow increases, fish screens, and revised gravel removal practices on Battle Creek (beginning summer 2006)
- Control of gravel mining, improvements of spawning areas, improvements of land management practices in the watershed, and

protection and restoration of riparian vegetation along Cottonwood Creek

Sacramento River National Wildlife Refuge

The Sacramento River National Wildlife Refuge (SRNWR) is composed of many units between the cities of Red Bluff and Princeton. The SRNWR along the middle Sacramento River is part of the Sacramento National Wildlife Refuge Complex, consisting of five refuges and three wildlife management areas within the Sacramento Valley. Reaches and subreaches of the river are delineated based generally on transitions in fluvial geomorphic riverine conditions, although county boundaries were considered as well. The middle Sacramento River region between Red Bluff and Colusa includes three units within the Chico Landing Subreach that contain restoration project sites addressed in the *Sacramento River–Chico Landing Subreach Habitat Restoration Draft Environmental Impact Report* (CBDA 2005). In addition, three areas proposed for restoration in this area occur within the larger SRNWR units that were evaluated in the *Environmental Assessment for Proposed Restoration Activities on the Sacramento River National Wildlife Refuge* (USFWS 2001; CBDA 2005).

In June 2005, USFWS issued the *Sacramento River National Wildlife Refuge Draft Comprehensive Conservation Plan and Environmental Assessment and Finding of No Significant Impact* (USFWS 2005) to serve as an integrated management plan for land that it acquires and manages for inclusion in the SRNWR. The SRNWR final comprehensive conservation plan includes goals, objectives, and strategies to guide management of lands within the SRNWR. It also includes assessments of and establishes parameters for “compatible uses,” which are uses that are considered compatible with the primary purposes for which the area was established. Riparian habitat restoration projects are being implemented under cooperative agreements between USFWS and other entities such as The Nature Conservancy (TNC) in accordance with the SRNWR final comprehensive conservation plan.

Sacramento River Preservation Trust

The Sacramento River Preservation Trust is a private, nonprofit organization active in environmental education and advocacy to preserve the natural environmental values of the Sacramento River. The trust has participated in various conservation and land acquisition projects, including securing lands for the SRNWR. The group is pursuing designation of a portion of the Sacramento River between Redding and Red Bluff as a national conservation area.

Sacramento River Watershed Program

The Sacramento River Watershed Program is an effort to bring stakeholders together to share information and work together to address water quality and other water-related issues within the Sacramento River watershed. The group is funded congressionally through the U.S. Environmental Protection Agency. The program’s primary goal is “to ensure that current and potential uses of

Sacramento River watershed resources are sustained, restored, and where possible, enhanced while promoting the long-term social and economic vitality of the region.” The Sacramento River Watershed Program manages grants for the Sacramento River Toxic Pollutants Control Program; performs extensive water quality monitoring, data collection, and data management for the watershed; and is instrumental in the study and monitoring of toxic pollutants. Although the program does not implement restoration projects, it is a potential partner for coordinating research and monitoring through consensus-based collaborative partnerships and promoting mutual education among the stakeholders of the Sacramento River watershed.

Sacramento Watersheds Action Group

The Sacramento Watersheds Action Group (SWAG) is a nonprofit corporation that secures funding for, designs, and implements projects that provide watershed restoration, streambank and slope stabilization, erosion control, watershed analysis, and road removal. SWAG has successfully worked with local groups, agencies, and organizations to fund and complete restoration projects on the Sacramento River and tributaries downstream from Keswick Dam. Their projects include development of the *Sulphur Creek Watershed Analysis and Action Plan*, the Whiskeytown Lake Shoreline Erosion Control Project, the Sulphur Creek Crossing Restoration Project, and the Lower Sulphur Creek Realignment and Riparian Habitat Enhancement Project. SWAG is a potential local sponsor for watershed restoration actions in the study area.

Shasta Land Trust

The Shasta Land Trust is a regional, nonprofit organization dedicated to conserving open space, wildlife habitat, and agricultural land. The trust works with public agencies and private landowners and is funded primarily through membership dues and donations. It employs various voluntary programs to protect and conserve valuable lands using conservation easements, land donations, and property acquisitions. The trust is a potential local partner for restoration activities in the Shasta Dam-to-Red Bluff area.

The Nature Conservancy

TNC is a private, nonprofit organization involved in environmental restoration and conservation throughout the United States and the world. TNC approaches environmental restoration primarily through strategic land acquisition from willing sellers and obtaining conservation easements. Some of the lands are retained by TNC for active restoration, research, or monitoring activities, while others are turned over to government agencies such as USFWS or DFG for long-term management. Lower in the Sacramento River basin, TNC has been instrumental in acquiring and restoring lands in the SRNWR and managing several properties along the Sacramento River. It also has pursued conservation easements on various properties at tributary confluences, including Cottonwood and Battle Creeks.

The Trust for Public Land

The Trust for Public Land is a national, nonprofit organization involved in preserving lands with natural, historic, cultural, or recreational value, primarily through conservation real estate. The trust's Western Rivers Program has been involved in conservation efforts along the Sacramento River between Redding and Red Bluff (the U.S. Bureau of Land Management's Sacramento River Bend Management Area), Battle Creek, Paynes Creek, Inks Creek, and Fenwood Ranch in Shasta County. The group promotes public ownership of conservation lands to ensure public access and enjoyment.

Chapter 2 Botanical Resources and Wetlands Attachments

This technical report includes the following attachments:

- Attachment 1, “Lists of All Special-Status Plant Species Known from or Potentially Present in the Primary and Extended Study Areas”
- Attachment 2, “List of Plant Species Observed in the Shasta Lake and Vicinity Portion of the Primary Study Area”
- Attachment 3, “Special-Status Plant Species Known to Occur in the Shasta Lake and Vicinity Portion of the Primary Study Area”
- Attachment 4, “List of All Sensitive Plant Species in the Extended Study Area Reported to the CNDDDB”
- Attachment 5, “Known Weed Source Locations, Potential Mode of Spread, and Risk of Spread”

This page left blank intentionally.

Chapter 3 Bibliography

- Atwater, B.F., and C.W. Hedel. 1976. Distribution of Seed Plants with Respect to Tide Levels and Water Salinity in the Natural Tidal Marshes of the Northern San Francisco Bay Estuary, California. Open-file Report 76-389. U.S. Geological Survey. Menlo Park, California.
- Bossard, C. C., J. M. Randall, and M. C. Hoshovsky. 2000. Invasive Plants of California's Wildlands. University of California Press. Berkeley, California.
- Burns, R.M., and B.H. Honkala. 1990. Silvics of North America: Volume 2, Hardwoods. Agricultural Handbook 654. U.S. Forest Service. Washington, D.C.
- CALFED. *See* CALFED Bay-Delta Program.
- CALFED Bay-Delta Program. 2000a (July). Multi-Species Conservation Strategy. Sacramento, California. Prepared by Jones & Stokes, Sacramento, California.
- . 2000b (August 28). Final Programmatic Environmental Impact Statement and Environmental Impact Report. Programmatic Record of Decision. Sacramento, California.
- . 2000c. Ecosystem Restoration Program Plan Volume I: Ecological Attributes of the San Francisco Bay-Delta Watershed. Final Programmatic Environmental Impact Statement/Environmental Impact Report Technical Appendix. Sacramento, California.
- California Bay-Delta Authority. 2005 (June). Sacramento River-Chico Landing Subreach Habitat Restoration Project Draft Environmental Impact Report. State Clearinghouse No. 2004112024. Sacramento, California. Lead agency: California Bay-Delta Authority. Project proponent: The Nature Conservancy. Prepared by EDAW, Sacramento, California. Available: <http://calwater.ca.gov/Programs/EcosystemRestoration/Ecosystem_Draft_ChicoLanding.shtml>.
- California Invasive Plant Council. 2006. California Invasive Plant Inventory. Cal-IPC Publication 2006-02. Berkeley, California.
- Cal-IPC. *See* California Invasive Plant Council.

- California Native Plant Society. 2007. Electronic Inventory of Rare and Endangered Vascular Plants of California. Available: <<http://cnps.web.aplus.net/cgi-bin/inv/inventory.cgi>>. Last updated July 9, 2007. Accessed August 29 and 30, 2007.
- . 2008. Electronic Inventory of Rare and Endangered Vascular Plants of California. Available: <<http://cnps.web.aplus.net/cgi-bin/inv/inventory.cgi>>. Accessed June 8, 2008.
- California Natural Diversity Database. 2007 (March). Rarefind: A Database Application for the Use of the California Department of Fish and Game's Natural Diversity Database. California Natural Heritage Division, California Department of Fish and Game, Sacramento, California.
- CBDA. *See* California Bay-Delta Authority.
- City of Colusa. 2007 (May). City of Colusa Draft General Plan. Colusa, California.
- City of Redding. 2000(October 30). City of Redding. 2000–2020 General Plan. Adopted by Resolution No. 2000-166. Redding, California.
- CNDDB. *See* California Natural Diversity Database.
- CNPS. *See* California Native Plant Society.
- Conard, S.G., R.L. MacDonald, and R.F. Holland. 1977. Riparian Vegetation and Flora of the Sacramento Valley. Pages 47–55 in A. Sands (ed.), *Riparian Forests of California: Their Ecology and Conservation*. Institute of Ecology Publication No. 15. University of California. Davis, California.
- Coops, H.F., W.B. Van Der Brink, and G. Van Der Velde. 1996. Growth and Morphological Responses of Four Helophyte Species in an Experimental Water-Depth Gradient. *Aquatic Botany* 54:11–24.
- Coops, H., R. Boeters, and H. Smith. 1991. Direct and Indirect Effects of Wave Attack on Helophytes. *Aquatic Botany* 41:333–352.
- DFG. *See* California Department of Fish and Game.
- DWR. *See* California Department of Water Resources.
- Environmental Laboratory. 1987. Corps of Engineers Wetlands Delineation Manual. U.S. Army Engineer Waterways Experiment Station. Vicksburg, Mississippi.

- Flora of North America Editorial Committee. 2002. Flora of North America North of Mexico Volume 23: Magnoliophyta: Commelinidae (in Part): Cyperaceae. Oxford University Press. New York, New York.
- Glenn County. 1993 (June 15). Glenn County General Plan. Volume I. Policies. Glenn County Planning and Public Works Agency. Willows, California.
- Griffin, J.R., and W.B. Critchfield. 1972. The Distribution of Forest Trees in California. Research Paper PSW-82, Pacific Southwest Forest and Range Experiment Station. U.S. Department of Agriculture. Berkeley, California.
- Hart, J.A., and J.C. Hunter. 2004. Restoring Slough and River Banks with Biotechnical Methods in the Sacramento–San Joaquin Delta. *Ecological Restoration* 22:262–268.
- Hickman, J.C. (ed.). 1993. The Jepson Manual: Higher Plants of California. University of California Press. Berkeley and Los Angeles, California.
- Holland, R.F. 1986. Preliminary Descriptions of the Terrestrial Natural Communities of California. Nongame-Heritage Program. Sacramento, California.
- Hunter, J.C., J.C. Sterling, W.P. Widdowson, E.C. Beedy, D. Stralberg, and N. Nur. 2003. The Abundance and Distribution of Non-native Woody Species in Sacramento Valley Riparian Zones. Pages 39-45 in C. Piroosko (ed.), *Proceedings California Invasive Plant Council Symposium Volume 7: 2003*. California Invasive Plant Council, Berkeley, California.
- Hunter, J.C., K.B. Willett, M.C. McCoy, J.F. Quinn, and K.E. Keller. 1999. Prospects for Preservation and Restoration of Riparian Forests in the Sacramento Valley, California, USA. *Environmental Management* 24(1):65–75.
- Jepson, W.L. 1893. The Riparian Botany of the Lower Sacramento. *Erythea* 1:238–246.
- . 1910. *Juglans californica* Wats (California black walnut). Pages 192–196 in the *Silva of California*. Memoirs of the University of California, Volume 2. The University Press. Berkeley, California.
- Keddy, P.A. 2000. *Wetland Ecology: Principles and Conservation*. Cambridge University Press. New York, New York.
- Lindstrand, L., III and J. Nelson. 2006. Habitat, Geologic, and Soil Characteristics of Shasta Snow-Wreath (*Neviusia cliftonii*) Populations. *Madrono* 53(1):65–68.

- Marty, J.T. 2005. Effects of Cattle Grazing on Diversity in Ephemeral Wetlands. *Conservation Biology* 19:1626–1632.
- Mason, H.L. 1957. *A Flora of the Marshes of California*. University of California Press. Berkeley, California.
- Mayer, K.E., and W.F. Laudenslayer (eds.). 1988. *A Guide to the Wildlife Habitats of California*. California Department of Forestry and Fire Protection. Sacramento, California.
- Nelson, J.R. 1994. Rare Plant Survey Guidelines. In M.W. Skinner and B.M. Pavlick (eds.), *Inventory of Rare and Endangered Vascular Plants of California*. California Native Plant Society. Sacramento, California.
- NRCS. *See* U.S. Natural Resources Conservation Service.
- Pyke, C.R. 2004. Simulating Vernal Pool Hydrologic Regimes for Two Locations in California, USA. *Ecological Modelling* 173:109–127.
- Reclamation. *See* U.S. Department of the Interior, Bureau of Reclamation.
- Reed, P.B., Jr. 1988. *National List of Plant Species that Occur in Wetlands: California (Region 0)*. U.S. Fish and Wildlife Service. Washington, D.C.
- Resources Agency. *See* The Resources Agency, State of California.
- RHJV. *See* Riparian Habitat Joint Venture.
- Riparian Habitat Joint Venture. 2004. *The Riparian Bird Conservation Plan: A Strategy for Reversing the Decline of Riparian Associated Birds in California*. Version 2.0. California Partners in Flight, Point Reyes Bird Observatory, Stinson Beach, California.
- Robins, J.D., and J.E. Vollmar. 2002. Livestock Grazing and Vernal Pools. Pages 401-430 in J. E. Vollmar (ed.), *Wildlife and Rare Plant Ecology of Eastern Merced County's Vernal Pool Grasslands*. Vollmar Consulting, Berkeley, California.
- Sacramento County. 1993 (December). *County of Sacramento General Plan Land Use Element*. Adopted December 15, 1993, and revised May 2, 1997, including Policy Amendments per the Agricultural-Residential Study adopted on February 11, 2004, by Resolution No. 2004-0174, Control No. 02-GPB-0379. Planning and Community Development Department. Sacramento, California.
- Sawyer, J.O., and T. Keeler-Wolf. 1995. *A Manual of California Vegetation*. California Native Plant Society. Sacramento, California.

- Sawyer, J.O., T. Keeler-Wolf, and J.M. Evens. 2009. *A Manual of California Vegetation* (2nd ed.). California Native Plant Society Press. Sacramento, California.
- Shasta County. 2004. *Shasta County General Plan*. Redding, California.
- Smith, R. E. 1912. *Walnut Culture in California: Walnut Blight*. Bulletin 231, Agricultural Experiment Station, College of Agriculture, University of California, Berkeley, California.
- Soil Survey Staff. *See* Soil Survey Staff, U.S. Natural Resources Conservation Service.
- Soil Survey Staff, U.S. Natural Resources Conservation Service. 2010. Web Soil Survey. Available: <<http://websoilsurvey.nrcs.usda.gov/>>.
- Solomeshch, A.I., M.G. Barbour, and R.F. Holland. 2007. Vernal Pools. Pages 394-424 in Barbour, M.G., T. Keeler-Wolf, and A.A. Schoenherr (eds.), *Terrestrial Vegetation of California*. University of California Press. Berkeley, California.
- The Resources Agency, State of California. 1989. *Upper Sacramento River Fisheries and Riparian Habitat Management Plan*. Prepared by SB 1086 Advisory Council.
- . 2003 (September). *Sacramento River Conservation Area Forum Handbook*. Sacramento, California. Prepared by the Sacramento River Advisory Council and revised and updated by the Sacramento River Conservation Area Forum.
- Thompson, K. 1961. Riparian Forests of the Sacramento Valley, California. *Annals of the Association of American Geographers* 51:294–315. Cited in Hunter et al. 1999.
- Tu, M. 2000. *Vegetation Patterns and Processes of Natural Regeneration in Periodically Flooded Riparian Forests in the Central Valley of California*. Dissertation. University of California, Davis, California.
- Tyler, C.M., B. Kuhn, and F.W. Davis. 2006. Demography and Recruitment Limitations of Three Oak Species in California. *Quarterly Review of Biology* 81(2):127–152.
- USACE. *See* U.S. Army Corps of Engineers.
- U.S. Army Corps of Engineers. 2006. *Interim Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Arid West Region*. U.S. Army Engineer Research and Development Center.

- U.S. Fish and Wildlife Service. 1995. Endangered and Threatened Wildlife and Plants; Proposed Endangered Status for Two Tidal Marsh Plants—*Cirsium hydrophilum* ssp. *hydrophilum* (Suisun thistle) and *Cordylanthus mollis* ssp. *mollis* (soft bird's beak)—from the San Francisco Bay Region. Federal Register 60(112):30999–31006 [Proposed Rules].
- . 2001 (May). Environmental Assessment for Proposed Restoration Activities on the Sacramento River National Wildlife Refuge (Ryan, Ohm, Haleakala, Pine Creek, Kaiser, Phelan Island, Koehnen, Hartley Island, and Stone Units). (J&S 00-003.) Willows, California. Prepared by Jones & Stokes, Sacramento, California.
- . 2005 (July). Sacramento River National Wildlife Refuge Final Comprehensive Conservation Plan and Environmental Assessment and Finding of No Significant Impact. Sacramento, California. Sacramento River Project. Chico, California.
- U.S. Forest Service. 1995a (April 28). Shasta-Trinity National Forest Land and Resource Management Plan. Shasta-Trinity National Forest, Redding, California.
- . 1995b. Forest Service Manual, Series 2000—National Forest Resource Management, WO amendment 2000-95-5, effective November 29, 1995.
- . 1996. Management Guide for the Shasta and Trinity Units of the Whiskeytown-Shasta-Trinity National Recreation Area. Redding, California.
- USFS. *See* U.S. Forest Service.
- USFWS. *See* U.S. Fish and Wildlife Service.
- U.S. Natural Resources Conservation Service. 2006. Field Indicators of Hydric Soils in the United States. Version 6.0. G.W. Hurt and L. M. Vasilas (eds.). U.S. Department of Agriculture. In cooperation with the National Technical Committee for Hydric Soils.
- Vaghti, M.G., and S.E. Greco. 2007. Riparian Vegetation of the Great Valley. Pages 425-449 in M. G. Barbour, T. Keeler-Wolf, and A. Schoenherr (eds.), *Terrestrial Vegetation of California*. University of California Press. Berkeley, California.